

AD \_\_\_\_\_

Award Number: W81XWH-07-1-0063

TITLE: Induction of the p75NTR by Aryl Propionic Acids in Prostate Cancer Cells

PRINCIPAL INVESTIGATOR: Shehla Wynne

CONTRACTING ORGANIZATION: Georgetown University  
Washington, DC 20057

REPORT DATE: December 2008

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 1 December 2008		2. REPORT TYPE Annual (Revised)		3. DATES COVERED 1 Dec 2007 – 30 Nov 2008	
4. TITLE AND SUBTITLE  Induction of the p75NTR by Aryl Propionic Acids in Prostate Cancer Cells				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-07-1-0063	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)  Shehla Wynne  E-Mail: sw254@georgetown.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Georgetown University Washington, DC 20057				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The p75 neurotrophin receptor (p75NTR) is a tumor suppressor in the prostate whose expression decreases as prostate cancer progresses. The purpose of this project is to investigate the role of p75NTR in the observed anticancer activity of aryl propionic acids in the prostate. It has been demonstrated that treatment of prostate cancer cells with the aryl propionic acids R-flurbiprofen and ibuprofen induces reexpression of p75NTR, decreases cell survival, and increases apoptosis. Investigation into the mechanism of p75NTR induction by R-flurbiprofen and ibuprofen revealed a strong correlation between increased p75NTR protein level and increased p75NTR mRNA level. Finally, induction of p75NTR seems to be dependent on the p38 MAPK pathway, which is involved in regulating mRNA stability of a subset of transcripts. Using the 2-phenyl propionic acid moiety of these profens as a pharmacophore, an in silico database of 30 million compounds was screened and carprofen was additionally identified as having activity for induction of p75NTR levels and inhibition of cell survival.					
15. SUBJECT TERMS p75NTR, ibuprofen, R-flurbiprofen, carprofen, prostate cancer, apoptosis, mRNA stability, p38 MAPK					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  UU	18. NUMBER OF PAGES  38	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

**Table of Contents**

	<u><b>Page</b></u>
<b>Introduction.....</b>	<b>4</b>
<b>Body.....</b>	<b>5</b>
<b>Key Research Accomplishments.....</b>	<b>6</b>
<b>Reportable Outcomes.....</b>	<b>8</b>
<b>Conclusions.....</b>	<b>9</b>
<b>References.....</b>	<b>11</b>
<b>Appendices.....</b>	<b>13</b>

## Introduction

The National Cancer Institute estimates there will be 218,890 new cases of prostate cancer and 27,050 deaths from prostate cancer in the United States this year (1). This makes prostate cancer the second most commonly diagnosed cancer type behind nonmelanoma skin cancer, and the second leading cause of cancer related death among men behind lung cancer (1). When detected early at the organ-confined stage, prostate cancer is often curable by surgical removal of the prostate or radiation therapy. However, when prostate cancer has spread outside of the prostate or has recurred following surgery or radiation therapy, current treatment involves some form of androgen deprivation therapy, and essentially always results in incurable hormone-refractory prostate cancer (HRPC). Once the cancer has reached this advanced stage, treatment options are limited and relatively unsuccessful. Docetaxel is the most successful drug to date and extends the life of the patient an average of two to three months. Therefore, there is a great need for improved treatment of advanced HRPC.

Nonsteroidal anti-inflammatory drugs (NSAIDs) demonstrate promise in both prevention and treatment of prostate cancer. They are traditionally used to treat inflammation by inhibiting cyclooxygenase (COX) activity. Increased COX-2 expression is believed to contribute to tumorigenesis through several mechanisms including stimulation of growth, promotion of angiogenesis, increased inflammation, increased invasion and migration, immune suppression, and inhibition of apoptosis (2). However the role of COX-2 in prostate cancer is controversial. Although, reports of COX-2 expression in prostate cancer vary, long term NSAID use is associated with decreased prostate cancer risk, and several COX inhibitors consistently induce apoptosis in prostate cancer cells regardless of COX-2 expression (3-7). Importantly, the efficacy of NSAIDs in inhibiting prostate cancer cell growth has been demonstrated in androgen nonresponsive cells such as PC-3 and DU-145, indicating potential for these drugs in the treatment of advanced prostate cancer (7-8). This is an exciting possibility given the limited options available for patients with HRPC. In particular, the aryl propionic acid class of NSAIDs, or profens, have repeatedly demonstrated anticancer activity in the prostate. This class of NSAIDs includes flurbiprofen, ibuprofen, naproxen, and ketoprofen among others. Long term ibuprofen use is associated with a decreased risk of prostate cancer (9-10). Treatment with the enantiomer R-flurbiprofen, which lacks COX inhibitory activity, was able to inhibit progression of prostate cancer in the TRAMP mouse (11). In addition, ibuprofen treatment decreased survival and induced apoptosis of DU-145 and LNCaP cells (7). This result was also observed in LNCaP cells treated with naproxen (7).

The p75 neurotrophin receptor (p75<sup>NTR</sup>) is an important player in the development of prostate cancer. It is a member of the tumor necrosis factor receptor superfamily (TNFR), and acts as a tumor suppressor in the prostate by inducing apoptosis and suppressing growth through its intracellular death domain (12-13). However, expression of p75<sup>NTR</sup> is decreased as prostate cancer progresses and is minimal in established advanced prostate cancer cell lines such as PC-3, DU-145, and LNCaP (14-15). Exogenous reexpression of p75<sup>NTR</sup> in prostate cancer cells resulted in decreased proliferation and increased apoptosis that was dependent upon the death domain of p75<sup>NTR</sup> (16). This indicates that drugs which induce reexpression of p75<sup>NTR</sup> in prostate cancer cells may have therapeutic potential. Interestingly, treatment of DU-145 prostate cancer cells with ibuprofen resulted in increased p75<sup>NTR</sup> expression (17). Therefore, it seems possible that induction of p75<sup>NTR</sup> may be causal of the observed anticancer activity of aryl propionic acids in the prostate.

## Body

**Task 1:** I took over this award as a continuation of the work of Ms Emily Quann, after the completion of her graduate degree. Previous studies supported by this award examined the effect of selected aryl propionic acid NSAIDs and structurally related compounds on the decreased survival of prostate cancer cell lines PC-3, DU-145, and LNCaP by induction of the p75<sup>NTR</sup> protein. The p75<sup>NTR</sup> has been shown to function as a tumor suppressor in the prostate by virtue of its intracellular death domain that can initiate apoptosis and inhibit growth. The most efficacious compounds for induction of p75<sup>NTR</sup> and decreased survival, in rank-order, were R-flurbiprofen, ibuprofen, oxaprozin, fenoprofen, naproxen, and ketoprofen. Since R-flurbiprofen and ibuprofen exhibited the greatest efficacy, these were the drugs used in subsequent studies. It was found that R-flurbiprofen and ibuprofen selectively induce p75<sup>NTR</sup>-dependent decreased survival of prostate cancer cells independently of COX inhibition. Previously our lab demonstrated that loss of p75<sup>NTR</sup> expression in prostate cancer cells may be due to increased p75<sup>NTR</sup> mRNA instability. Consistently, it was found that the observed increase in p75<sup>NTR</sup> protein due to R-flurbiprofen and ibuprofen treatment was accompanied by an increase in p75<sup>NTR</sup> mRNA, and this increase in mRNA was the result of increased p75<sup>NTR</sup> mRNA stability. In addition, treatment with R-flurbiprofen or ibuprofen led to sustained activation of the p38 MAPK pathway and inhibition of this pathway prevented an induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen. Collectively, the data suggest that R-flurbiprofen and ibuprofen induce p75<sup>NTR</sup> expression leading to p75<sup>NTR</sup> dependent decreased survival by increased p75<sup>NTR</sup> mRNA stability that is mediated through the p38 MAPK pathway.

Following Dr Quann's work, using the 2-phenyl propionic acid moiety of these profens as a pharmacophore, an in silico database of 30 million compounds was screened and carprofen was additionally identified as having activity for induction of p75<sup>NTR</sup> levels and inhibition of cell survival.

All procedures and results are described in detail in the manuscripts found in the appendices.

**Task 2:** The goal of task 2 is to investigate the role of the aryl hydrocarbon receptor (AhR) in the induction of p75<sup>NTR</sup> expression by R-flurbiprofen or ibuprofen. Our investigation of AhR identified a role of the p38 MAPK in the induction of p75<sup>NTR</sup>. I specifically examined the effect of inhibiting Dual Specificity Phosphatases, or DUSPs on the induction of in the prostate cancer cell lines PC-3, DU-145, and LNCaP. I scanned the cell lines for expression of Wip-1, HePTP, PP2C $\alpha$ , MKP-1, MKP-5, M3/6 and found that they are expressed at varying levels in the three cell lines. However it was found that inhibition of the DUSPs had little to no effect on the induction of p75<sup>NTR</sup>. Presently, my focus is on the two other MAPK pathways - the ERK and JNK MAPK pathways – to determine their involvement in the induction of p75<sup>NTR</sup> by profens. In addition I am investigating the role of NSAID mediated p75<sup>NTR</sup> induction in the cell cycle. This task is still under progress.

## Key Research Accomplishments

- Using the 2-phenyl propionic acid moiety of these profens as a pharmacophore, *in silico* database of 30 million compounds was screened. Carprofen as having an order of magnitude greater activity for induction of p75<sup>NTR</sup> levels and inhibition of cell survival (figure 1, Appendix 1).
- Prostate (PC-3 and DU-145) and bladder (T24) cancer cells were found to be more sensitive to carprofen induction of p75<sup>NTR</sup>-associated loss of survival than breast (MCF-7) and fibroblast (3T3) cells (figure 2 A and B, Appendix 1).
- Transfection of prostate cell lines with a dominant-negative form of p75<sup>NTR</sup> before carprofen treatment partially rescued cell survival, showing a cause-and-effect relationship between carprofen induction of p75<sup>NTR</sup> levels and inhibition of survival (figure 3 A and B, Appendix 1).
- Carprofen induced apoptotic nuclear fragmentation in the prostate cancer cell lines DU-145 and PC-3, but not in MCF-7 and 3T3 cells. Furthermore, small interfering RNA knockdown of the p38 mitogen-activated protein kinase (MAPK) protein prevented induction of p75<sup>NTR</sup> by carprofen in both prostate cancer cell lines (figure 4, Appendix 1).
- Carprofen treatment induced phosphorylation of p38 MAPK as early as within 1 min. Expression of a dominant-negative form of MK2, the kinase downstream of p38 MAPK frequently associated with signaling cascades leading to apoptosis, prevented carprofen induction of the p75<sup>NTR</sup> protein (figure 5 and figure 6, Appendix 1).

The items listed in the Statement of Work Task 1 below have been completed by the previous PI Dr Emily Quann. Some additional work was done and has been added to the accomplishments of Task 1. Task 2 has not yet been completed. These and other research accomplishments have been published in the following three journal articles and are included as Appendices.

Khwaja FS, Quann EJ, Pattabiraman N, Wynne S, Djakiew D. Carprofen induction of p75<sup>NTR</sup>-dependent apoptosis via the p38 mitogen-activated protein kinase pathway in prostate cancer cells. *Mol Cancer Ther.* 2008 Nov;7(11):3539-45.

Quann EJ, Khwaja F, Djakiew D. The p38 MAPK Pathway Mediates Aryl Propionic Acid-Induced Messenger RNA Stability of p75<sup>NTR</sup> in Prostate Cancer Cells. *Cancer Res* 2007;67:11402-10.

Quann EJ, Khwaja F, Zavitz KH, Djakiew D. The Aryl Propionic Acid R-Flurbiprofen Selectively Induces p75<sup>NTR</sup>-Dependent Decreased Survival of Prostate Tumor Cells. *Cancer Res* 2007;67:3254-62.

**Task 1:** Determine the role of p75<sup>NTR</sup> induction in aryl propionic acid induced survival inhibition of prostate cancer cells (1-18 months).

- Perform western blots for cyclooxygenase (COX) expression following 48 hour aryl propionic acid treatment of PC-3 or DU-145 cells.
- Perform western blots for expression of other tumor necrosis factor receptor (TNFR) superfamily members following 48 hour aryl propionic acid treatment of PC-3 or DU-145 cells.
- Perform western blots for p75<sup>NTR</sup> following 2 week chronic aryl propionic acid treatment of PC-3 or DU-145 cells at concentrations lower than those used in the 48 hour treatments.
- Determine enumeration of PC-3 and DU-145 cells using a hemocytometer following 2 week chronic aryl propionic acid treatment at concentrations lower than those used in the 48 hour treatments.
- Perform rescue experiments with PC-3 and DU-145 cells involving transfection of ecdysone-inducible dominant negative p75<sup>NTR</sup> expression vectors followed by 48 hour aryl propionic acid treatment. Determine relative survival by MTT assay.
- Perform rescue experiments with PC-3 and DU-145 cells involving 24 hour transfection of p75<sup>NTR</sup>-targeted siRNA followed by 48 hour aryl propionic acid treatment. Determine relative cell survival by MTT assay.
- The above listed experiments were repeated with the NSAID carprofen, and it was found to have an order of magnitude greater activity for the induction of p75<sup>NTR</sup>.

**Task 2:** Determine the role of proteins proximal to p38 MAPK in the induction of p75<sup>NTR</sup>.

- Perform western blots for expression of p75<sup>NTR</sup> in PC-3, Du-145 and LNCaP cells following 72 hour Dual Specificity Phosphatase targeted siRNA transfection.
- Perform western blots for expression of p75<sup>NTR</sup> in PC-3, DU-145 and LNCaP cells following 24 hour Dual Specificity Phosphatase dominant negative mutant transfection.
- Perform western blots for expression of p75<sup>NTR</sup> following 24 hours treatment with commercially available inhibitors for the ERK and JNK MAPK pathways.

## Reportable Outcomes

### Manuscripts

- Khwaja FS, Quann EJ, Pattabiraman N, Wynne S, Djakiew D. Carprofen induction of p75<sup>NTR</sup>-dependent apoptosis via the p38 mitogen-activated protein kinase pathway in prostate cancer cells. *Mol Cancer Ther*. 2008 Nov;7(11):3539-45.
- Quann EJ, Khwaja F, Zavitz KH, Djakiew D. The Aryl Propionic Acid R-Flurbiprofen Selectively Induces p75<sup>NTR</sup>-Dependent Decreased Survival of Prostate Tumor Cells. *Cancer Res* 2007;67:3254-62.
- Quann EJ, Khwaja F, Djakiew D. The p38 MAPK Pathway Mediates Aryl Propionic-Acid Induced Messenger RNA Stability of p75<sup>NTR</sup> in Prostate Cancer Cells. *Cancer Res* 2007;67:11402-10.

### Conference Abstracts

- Quann EJ, Khwaja F, Djakiew D. The p38 MAPK pathway mediates aryl propionic acid induced mRNA stability of p75<sup>NTR</sup> in prostate cancer cells. AACR-NCI-EORTC Molecular Targets and Cancer Therapeutics, October 2007, San Francisco.



## Conclusions

This study revealed that several aryl propionic acids induce reexpression of p75<sup>NTR</sup> in two different metastatic hormone-refractory prostate cancer cell lines, PC-3 and DU-145. Of those tested, the enantiomer R-flurbiprofen and ibuprofen were the most effective. These drugs were also effective in inducing p75<sup>NTR</sup> expression in the metastatic androgen responsive LNCaP cell line. R-flurbiprofen or ibuprofen treatment of all three cell lines resulted in decreased survival that corresponded with induction of p75<sup>NTR</sup>, confirming the potential for these drugs as anticancer agents in the prostate and also suggesting that p75<sup>NTR</sup> induction is involved in the observed decreased survival. Additionally, the drug carprofen was also shown to have activity for the induction of p75<sup>NTR</sup>. In addition, treatment with these drugs resulted in induction of apoptosis. Consistently, p75<sup>NTR</sup> is a TNFR family member capable of inducing apoptosis through its conserved intracellular death domain. There are several other TNFR family members that also possess apoptosis inducing death domains; however, none of these family members was significantly upregulated in response to profen treatment. Increases in p75<sup>NTR</sup> protein expression were found to closely correlate with increases in p75<sup>NTR</sup> mRNA following profen treatment. The observed increase in p75<sup>NTR</sup> mRNA level was found to be largely due to substantial increases in mRNA stability. The p38 MAPK pathway has been shown to regulate mRNA stability through its downstream kinases MK2 and MK3, and therefore, its role in p75<sup>NTR</sup> induction by profens was investigated. Profen treatment resulted in increased p38 MAPK phosphorylation, and inhibition of this pathway using a p38 MAPK inhibitor or by siRNA knockdown of p38 MAPK, or MK2 and MK3 before profen treatment prevented an induction of p75<sup>NTR</sup>. Collectively, the data suggests that profen treatment activates the p38 MAPK pathway, which increases the stability of the p75<sup>NTR</sup> transcript, thus resulting in increased p75<sup>NTR</sup> mRNA, increased p75<sup>NTR</sup> protein expression, and p75<sup>NTR</sup> mediated decreased prostate cancer cell survival.

Investigation into the role of p38 MAPK proximal proteins, specifically the Dual Specificity Phosphatases revealed that although expressed at varying levels, these phosphatases are not involved in the profen mediated induction of p75<sup>NTR</sup>. The DUSPs scanned were Wip-1, HePTP, PP2C $\alpha$ , MKP-1, MKP-5, M3/6 and PAC-1.

The results described above have several implications. They contribute to the body of evidence demonstrating the potential of aryl propionic acids for the treatment of prostate cancer, and identify induction of p75<sup>NTR</sup> as a COX independent mechanism by which these drugs achieve their anticancer activity in the prostate. This study also indicates that drugs which result in the reexpression of p75<sup>NTR</sup> may be effective in the treatment of advanced prostate cancer. It is significant that the profens were effective in inducing p75<sup>NTR</sup> expression and in decreasing survival of three different advanced prostate cancer cell lines, each derived from metastasis to a different organ, as this demonstrates that the results are not an artifact of a single cell line. This is an important observation given the significant dedifferentiation and heterogeneity associated with the advanced disease. In addition, these results were observed in both androgen independent and androgen responsive cells lines, suggesting these drugs may be effective for advanced prostate cancer regardless of AR status. Finally, this brings to light novel approaches for identifying drugs which induce p75<sup>NTR</sup> with greater specificity. For example, identification of drugs that activate p38 MAPK may reveal drugs that induce p75<sup>NTR</sup>. Better yet would be the identification of drugs which specifically activate the downstream kinases MK2 and MK3, as these may have decreased side effects relative to drugs that activate p38 MAPK which has many downstream targets. The exact mechanisms by which MK2 and MK3 stabilize mRNAs need to

be explored further. It is highly likely that several RNA binding proteins are involved in increasing p75<sup>NTR</sup> mRNA stability due to profen treatment, and further elucidation of this mechanism may lead to identification of additional novel targets and the identification and development of therapeutics highly specific in inducing p75<sup>NTR</sup> in prostate cancer cells with minimal off target effects.

## References

- 1) Comprehensive Cancer Information - National Cancer Institute. National Cancer Institute 2007. Internet: <http://www.cancer.gov/>.
- 2) Liao Z, Mason KA, Milas L. Cyclo-oxygenase-2 and its inhibition in cancer: is there a role? *Drugs* 2007;67:821-45.
- 3) Jacobs EJ, Thun MJ, Bain EB, Rodriguez C, Henley SJ, Calle EE. A large cohort study of long-term daily use of adult-strength aspirin and cancer incidence. *J Natl Cancer Inst* 2007;99:608-15.
- 4) Roberts RO, Jacobson DJ, Girman CJ, Rhodes T, Lieber MM, Jacobsen SJ. A population-based study of daily nonsteroidal anti-inflammatory drug use and prostate cancer. *Mayo Clin Proc* 2002;77:219-25.
- 5) Hsu AL, Ching TT, Wang DS, Song X, Rangnekar VM, Chen CS. The cyclooxygenase-2 inhibitor celecoxib induces apoptosis by blocking Akt activation in human prostate cancer cells independently of Bcl-2. *J Biol Chem* 2000;275:11397-403.
- 6) Johnson AJ, Song X, Hsu A, Chen C. Apoptosis signaling pathways mediated by cyclooxygenase-2 inhibitors in prostate cancer cells. *Adv Enzyme Regul* 2001;41:221-35.
- 7) Andrews J, Djakiew D, Krygier S, Andrews P. Superior effectiveness of ibuprofen compared with other NSAIDs for reducing the survival of human prostate cancer cells. *Cancer Chemother Pharmacol* 2002;50:277-84.
- 8) Kamiyo T, Sato T, Nagatomi Y, Kitamura T. Induction of apoptosis by cyclooxygenase-2 inhibitors in prostate cancer cell lines. *Int J Urol* 2001;8:S35-S39.
- 9) Jacobs EJ, Rodriguez C, Mondul AM et al. A large cohort study of aspirin and other nonsteroidal anti-inflammatory drugs and prostate cancer incidence. *J Natl Cancer Inst* 2005;97:975-80.
- 10) Nelson JE, Harris RE. Inverse association of prostate cancer and non-steroidal anti-inflammatory drugs (NSAIDs): results of a case-control study. *Oncol Rep* 2000;7:169-70.
- 11) Wechter WJ, Leipold DD, Murray ED, Jr. et al. E-7869 (R-flurbiprofen) inhibits progression of prostate cancer in the TRAMP mouse. *Cancer Res* 2000;60:2203-8.
- 12) Krygier S, Djakiew D. The neurotrophin receptor p75<sup>NTR</sup> is a tumor suppressor in human prostate cancer. *Anticancer Res* 2001;21:3749-55.
- 13) Allen J, Khwaja F, Djakiew D. Gene therapy of prostate xenograft tumors with a p75<sup>NTR</sup> lipoplex. *Anticancer Res* 2004;24:2997-3003.

- 14) Perez M, Regan T, Pflug B, Lynch J, Djakiew D. Loss of low-affinity nerve growth factor receptor during malignant transformation of the human prostate. *Prostate* 1997;30:274-9.
- 15) Krygier S, Djakiew D. Molecular characterization of the loss of p75<sup>NTR</sup> expression in human prostate tumor cells. *Mol Carcinog* 2001;31:46-55.
- 16) Allen J, Khwaja F, Byers S, Djakiew D. The p75<sup>NTR</sup> mediates a bifurcated signal transduction cascade through the NF kappa B and JNK pathways to inhibit cell survival. *Exp Cell Res* 2005;304:69-80.
- 17) Khwaja F, Allen J, Lynch J, Andrews P, Djakiew D. Ibuprofen inhibits survival of bladder cancer cells by induced expression of the p75<sup>NTR</sup> tumor suppressor protein. *Cancer Res* 2004;64:6207-13.

## Appendices

Khwaja FS, Quann EJ, Pattabiraman N, Wynne S, Djakiew D. Carprofen induction of p75<sup>NTR</sup>-dependent apoptosis via the p38 mitogen-activated protein kinase pathway in prostate cancer cells. *Mol Cancer Ther*. 2008 Nov;7(11):3539-45.

Quann EJ, Khwaja F, Zavitz KH, Djakiew D. The Aryl Propionic Acid R-Flurbiprofen Selectively Induces p75<sup>NTR</sup>-Dependent Decreased Survival of Prostate Tumor Cells. *Cancer Res* 2007;67:3254-62.

Quann EJ, Khwaja F, Djakiew D. The p38 MAPK Pathway Mediates Aryl Propionic Acid-Induced Messenger RNA Stability of p75<sup>NTR</sup> in Prostate Cancer Cells. *Cancer Res* 2007;67:11402-10.

# Carprofen induction of p75<sup>NTR</sup>-dependent apoptosis via the p38 mitogen-activated protein kinase pathway in prostate cancer cells

Fatima S. Khwaja,<sup>1</sup> Emily J. Quann,<sup>3</sup>  
Nagarajan Pattabiraman,<sup>2</sup> Shehla Wynne,<sup>1</sup>  
and Daniel Djakiew<sup>1,2</sup>

<sup>1</sup>Department of Biochemistry and Molecular & Cellular Biology and <sup>2</sup>Vincent T. Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, District of Columbia; and <sup>3</sup>Immunology Program, Memorial Sloan Kettering Cancer Center, New York, New York

## Abstract

The p75 neurotrophin receptor (p75<sup>NTR</sup>) functions as a tumor suppressor in prostate epithelial cells, where its expression declines with progression to malignant cancer. Previously, we showed that treatment with *R*-flurbiprofen or ibuprofen induced p75<sup>NTR</sup> expression in several prostate cancer cell lines leading to p75<sup>NTR</sup>-mediated decreased survival. Using the 2-phenyl propionic acid moiety of these profens as a pharmacophore, we screened an *in silico* database of 30 million compounds and identified carprofen as having an order of magnitude greater activity for induction of p75<sup>NTR</sup> levels and inhibition of cell survival. Prostate (PC-3 and DU-145) and bladder (T24) cancer cells were more sensitive to carprofen induction of p75<sup>NTR</sup>-associated loss of survival than breast (MCF-7) and fibroblast (3T3) cells. Transfection of prostate cell lines with a dominant-negative form of p75<sup>NTR</sup> before carprofen treatment partially rescued cell survival, showing a cause-and-effect relationship between carprofen induction of p75<sup>NTR</sup> levels and inhibition of survival. Carprofen induced apoptotic nuclear fragmentation in prostate but not in MCF-7 and 3T3 cells. Furthermore, small interfering RNA knockdown of the p38 mitogen-activated protein kinase (MAPK) protein prevented induction of p75<sup>NTR</sup> by carprofen in both prostate cell lines. Carprofen treatment induced phosphorylation of p38 MAPK as early as within 1 min.

Expression of a dominant-negative form of MK2, the kinase downstream of p38 MAPK frequently associated with signaling cascades leading to apoptosis, prevented carprofen induction of the p75<sup>NTR</sup> protein. Collectively, we identify carprofen as a highly potent profen capable of inducing p75<sup>NTR</sup>-dependent apoptosis via the p38 MAPK pathway in prostate cancer cells. [Mol Cancer Ther 2008;7(11):3539–45]

## Introduction

The p75 neurotrophin receptor (p75<sup>NTR</sup>) is a 75-kDa cell surface receptor glycoprotein that shares both structural and sequence homology with the tumor necrosis factor receptor superfamily of proteins (1, 2). Some of these proteins (e.g., p75<sup>NTR</sup>, p55<sup>TNFR</sup>, Fas, DRs3-6, and EDAR) have similar sequence motifs of defined elongated structure (1) designated “death domains” based on their apoptosis-inducing function (2). In the human prostate, the p75<sup>NTR</sup> protein is progressively lost in pathologic cancer tissues (3). The proportion of epithelial cells that have retained p75<sup>NTR</sup> expression in the organ-confined pathologic prostate is inversely associated with increasing Gleason score and preoperative serum prostate-specific antigen concentrations (4). In addition, immunoblot of human prostate epithelial cell lines derived from metastases exhibit a further reduction of p75<sup>NTR</sup> expression (5). Significantly, although expression of the p75<sup>NTR</sup> protein is suppressed, the gene encoding p75<sup>NTR</sup> appears intact in these prostate cancer cells (6). The loss of p75<sup>NTR</sup> expression is a result of a loss of mRNA stability (6). Following ectopic reexpression of the p75<sup>NTR</sup> in these cancer cells, their rate of apoptosis increased (7). Additionally, the same ectopically expressing p75<sup>NTR</sup> cancer cells exhibited a retardation of cell cycle progression characterized by accumulation of cells in G<sub>1</sub> phase with a corresponding reduction of cells in the S phase of the cell cycle (7). Consistent with these observations, the p75<sup>NTR</sup> has been characterized with both tumor suppressor and metastasis suppressor activity in prostate cancer cells (7, 8).

Several studies have shown that nonsteroidal anti-inflammatory drugs (NSAID) are effective as anticancer agents for colorectal, breast, pancreatic, squamous cell carcinoma of the head and neck, bladder, ovarian, lung, and prostate cancers (9, 10). With respect to prostate cancer, retrospective studies indicate that there is a significantly reduced risk of prostate cancer associated with regular use of NSAIDs (11–13). *In vivo* studies using rodents have indicated that NSAIDs can decrease the size of prostate tumors (14, 15) and suppress the metastasis of prostatic cancer (14, 16). There is no common mechanism of action underlying NSAIDs effectiveness against cancer cells. Some

Received 6/2/08; revised 8/13/08; accepted 8/24/08.

**Grant support:** NIH grants R01DK52626 and U56CA101429 and Department of Defense Prostate Cancer Research Program grant PC060409.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Requests for reprints:** Daniel Djakiew, Department of Biochemistry and Molecular & Cellular Biology, Georgetown University Medical Center, 3900 Reservoir Road Northwest, Washington, DC 20057-1436. Phone: 202-687-1203; Fax: 202-687-1823. E-mail: djakiewd@georgetown.edu

Copyright © 2008 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-08-0512


NSAIDs inhibit the cyclooxygenases (COX) that convert arachidonic acid to prostaglandins (17). Prostaglandins are thought to contribute to tumor growth by inhibiting apoptosis (18) and by inducing the formation of new blood vessels needed to sustain tumor growth (19). Hence, COX inhibition of prostaglandin synthesis could explain part of the antitumor activity of certain NSAIDs. However, NSAIDs can also inhibit tumor formation and growth of COX-null cell lines (20). In addition, NSAIDs that lack COX inhibitory activity can still have significant anticancer effects both *in vivo* (21) and *in vitro* (22). Similarly, growth of the DU-145 prostate cancer cell line that lacks expression of COX-1 and COX-2 is inhibited by NSAIDs (23). Interestingly, *R*-flurbiprofen and ibuprofen have been shown to induce p75<sup>NTR</sup> levels leading to apoptosis in prostate cancer cell lines (23). These profens activated the p38 mitogen-activated protein kinase (MAPK) pathway leading to stabilization of p75<sup>NTR</sup> mRNA and increased levels of p75<sup>NTR</sup> protein that subsequently induced apoptosis of the prostate cancer cells (24). In this report, we used the 2-phenyl propionic acid moiety of the profens

as a pharmacophore for an *in silico* search of related compounds and identified carprofen as having an order of magnitude greater activity for induction of p75<sup>NTR</sup> levels and inhibition of cell survival. Carprofen activity occurred through rapid phosphorylation of p38 MAPK, which signaled through MK2 to increase levels of p75<sup>NTR</sup> protein and stimulate apoptosis in the prostate cancer cells.

## Materials and Methods


### Cell Lines and Culture Conditions

PC-3 and DU-145 prostate cell lines were obtained from the Tissue Culture Core Facility of the Georgetown University Lombardi Comprehensive Cancer Center. T24 bladder, MCF-7 breast, and 3T3 fibroblast cells were obtained from the American Type Culture Collection. All cell lines were maintained in DMEM (Mediatech) containing 4.5g/L glucose and L-glutamine supplemented with antibiotic/antimycotic [100 units/mL penicillin G, 100 µg/mL streptomycin, and 0.25 µg/mL amphotericin B (Mediatech)] and 10% fetal bovine serum (Sigma). Cells were incubated in the presence of 5% CO<sub>2</sub> and air at 37°C.



**Figure 1.** Immunoblots of p75<sup>NTR</sup> levels in PC-3 and DU-145 cells following 48 h treatment with 0, 0.1, 0.25, 0.5, and 1.0 mmol/L drug. The compound ID of each drug is given adjacent to its chemical structure. The A875 melanoma cell line was used as a positive control for p75<sup>NTR</sup> expression.

**Figure 2. A**, immunoblots of p75<sup>NTR</sup> levels with corresponding  $\beta$ -actin loading controls in PC-3 and DU-145 prostate cancer cells, T24 bladder cancer cells, MCF-7 breast cancer cells, and 3T3 fibroblasts after 48 h treatment with 0, 20, 40, 60, 80, and 100  $\mu$ M carprofen. A875 cell lysates were used as positive controls for p75<sup>NTR</sup> expression. **B**, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide cell survival assay of PC-3 and DU-145 prostate cancer cells, T24 bladder cancer cells, MCF-7 breast cancer cells, and 3T3 fibroblasts following 48 h treatment with 0, 20, 40, 60, 80, and 100  $\mu$ M carprofen. \*,  $P < 0.01$ .



### Drug Preparation, Treatment, and Cell Lysis

Using 2-phenyl propionic acid as a pharmacophore, we searched an *in silico* database of ~30 million compounds from which nine aryl propionic acids were selected for further analysis. Stock solutions were prepared by dissolving each aryl propionic acid in DMSO (Sigma) at a concentration of 200 mmol/L. Cells were seeded overnight at 70% to 80% confluency and were then treated for 48 h at concentrations of 0, 20, 40, 60, 80, and 100  $\mu$ M/L. Cell lysates of treated cells were prepared as described previously (7, 8, 23). The supernatant was retained and protein concentration was determined according to the manufacturer's protocol (Bio-Rad Laboratories).

### Immunoblot Analysis

Immunoblot analysis was done as described previously (23). Membranes were incubated in the primary antibody: murine monoclonal anti-p75<sup>NTR</sup> (1:2,000; Upstate Cell Signaling Solutions), rabbit polyclonal phosphorylated p38 MAPK (1:1,000), mouse monoclonal anti-p38 $\alpha$  (1:1,000; Cell Signaling Technology), or murine monoclonal anti- $\beta$ -actin (1:5,000; Sigma). Membranes were subsequently incubated in goat anti-mouse or goat anti-rabbit horseradish peroxidase-conjugated secondary antibodies (Bio-Rad Laboratories) at a dilution of 1:2,000 and immunoreactivity was visualized with a chemiluminescence detection reagent (Amersham Pharmacia Biotech). The positive control for p75<sup>NTR</sup> expression was a whole-cell lysate of A875 cells (Dr. Moses Chao, Cornell University).

### Cell Survival Assay with p75<sup>NTR</sup> Dominant-Negative Transfection and Hoechst Dye Nuclear Staining

An equal number of viable cells ( $2 \times 10^3$  per well) in 96-well culture plates (final volume of 100  $\mu$ L/well culture medium) were incubated for 4 h at 37°C in a humidified atmosphere of 5% CO<sub>2</sub>. Some cells were also transiently transfected with a p75<sup>NTR</sup> dominant-negative vector described previously (24–26). The  $\Delta$ ICD vector expresses a p75<sup>NTR</sup> gene product with the intracellular domain (ICD) deleted. The  $\Delta$ ICD is an ecdysone-inducible p75<sup>NTR</sup> vector and therefore was cotransfected with the ecdysone receptor plasmid pVgRxR. The transfection was done with Lipofect-

AMINE reagent (Invitrogen) in serum-free medium for 6 h, after which serum-containing medium was added. After 18 subsequent hours, cells were incubated in 1  $\mu$ M/L ponasterone A (Invitrogen) for 24 h to drive expression of the dominant-negative gene product. Following incubation with ponasterone A, cells were treated with carprofen (0–100  $\mu$ M/L) for 48 h and relative cell survival was determined with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide labeling reagent (final concentration, 0.5 mg/mL; Roche Diagnostics). Subsequently, cells were incubated overnight with 100  $\mu$ L/well solubilization solution and the samples were quantified at 570 nm using a microtiter plate reader (Bio-Rad Laboratories). Hoechst dye nuclear (DNA) staining to identify apoptotic nuclei was conducted as described previously (25). PC-3, DU-145, MCF-7, and 3T3 cells were treated for 48 h with carprofen and then fixed in 10% formalin (Electron Microscopy Sciences). Some cells were transfected with the  $\Delta$ ICDp75<sup>NTR</sup> plus ponasterone A before carprofen treatment.

### Small Interfering RNA Transfection

Cells were transfected for 72 h with nontargeting small interfering RNA (siRNA) or siRNA specific for p38 $\alpha$  (J-003512-20; Dharmacon RNA Technologies) at final concentrations of 100 nmol/L according to the manufacturer's protocol. Transfection reagent DharmaFECT 1 was used for DU-145 cells, and DharmaFECT 2 was used for PC-3 cells (Dharmacon RNA Technologies). After transfection, the cells were treated with carprofen for 48 h followed by determination of p75<sup>NTR</sup> protein expression.

### MK2 Dominant-Negative Transfection

PC-3 and DU-145 cells were transiently transfected with a MK2 dominant-negative vector (MK2-K76R) described previously (27). The transfection was done with Lipofect-AMINE reagent (Invitrogen) in serum-free medium for 6 h, after which serum-containing medium was added for 24 h to allow expression of the dominant-negative gene product. Cells were treated with carprofen (100  $\mu$ M/L) for 48 h and expression of p75<sup>NTR</sup> protein was determined by immunoblot with mouse monoclonal anti-p75<sup>NTR</sup> (1:2,000; Millipore).



### Statistical Analysis

The statistical differences between data sets and/or means were analyzed by ANOVA or the Mann-Whitney test using the Prism program (GraphPad Software) and the data expressed as the mean  $\pm$  SE. Data were considered statistically significant when  $P \leq 0.05$ .

## Results

### Carprofen Exhibits Superior Efficacy of the Aryl Propionic Acids to Induce p75<sup>NTR</sup> Levels Associated with Cell-Specific Decreased Survival


Analysis of the 2-phenyl propionic acid pharmacophore homology search identified nine aryl propionic acids that were screened for activity to induce expression of p75<sup>NTR</sup> protein in PC-3 and DU-145 human prostate cancer cells. Initially, the PC-3 and DU-145 cell lines were selected because they are the only two prostate tumor cell lines included in the NIH Developmental Therapeutics Program Anticancer Drug Discovery Program. The immunoblots showing activity of each compound to induce p75<sup>NTR</sup> were placed in rank-order (Fig. 1). In both cell

lines, carprofen exhibited superior efficacy for induction of p75<sup>NTR</sup> expression at a concentration of  $\leq 100$   $\mu\text{mol/L}$  compared with all other aryl propionic acids examined (Fig. 1). At lower concentrations, carprofen selectively induced expression of p75<sup>NTR</sup> protein at  $\geq 40$   $\mu\text{mol/L}$  in PC-3 and DU-145 prostate cancer cells, as well as in the T24 bladder cancer cell line, but not in the MCF-7 breast cancer cell line or the 3T3 fibroblast cell line (Fig. 2A). The T24 bladder cancer cell line was included as a positive control because they were shown previously to be sensitive to profen (ibuprofen and *R*-flurbiprofen)-induced p75<sup>NTR</sup>-dependent decreased survival, whereas MCF-7 and 3T3 cells were included as negative controls because they were shown previously not to be sensitive to profen (ibuprofen and *R*-flurbiprofen)-induced decreased survival (25).

Carprofen treatment selectively decreased the survival of cells in rank-order with PC-3 and DU-145 prostate cancer cells exhibiting greatest sensitivity to dose-dependent decreased survival followed by the T24 bladder cancer cells and with MCF-7 and 3T3 fibroblasts the least sensitive to carprofen-induced decreased survival (Fig. 2B). Significantly, there was a strong association between the dose-dependent induction of p75<sup>NTR</sup> levels (Fig. 2A) and decreased survival of specific cell types following carprofen treatment (Fig. 2B).


### Carprofen Induced Decreased Prostate Cancer Cell Survival Is Dependent on p75<sup>NTR</sup>

To establish a causal relationship between carprofen induction of p75<sup>NTR</sup> protein expression and inhibition of cell survival, we used a ponasterone A-inducible expression vector for p75<sup>NTR</sup> that exhibits a deletion of the intracellular death domain ( $\Delta\text{ICDp75}^{\text{NTR}}$ ) shown to function as a dominant-negative antagonist of the intact p75<sup>NTR</sup> gene product (23–26). The treatment of both PC-3 and DU-145 cells with carprofen or carprofen plus ponasterone A inhibited cell survival in a dose-dependent manner (Fig. 3). However, both PC-3 and DU-145 cell lines induced with ponasterone A to express  $\Delta\text{ICDp75}^{\text{NTR}}$  exhibited a significant ( $P < 0.001$ ) partial rescue from carprofen-mediated inhibition of cell survival relative to carprofen-treated  $\Delta\text{ICDp75}^{\text{NTR}}$  cells in the absence of ponasterone A (Fig. 3). Subsequently, we examined Hoechst-stained nuclear morphology to identify fragmented nuclei typical of apoptotic cells with the exception of T24 bladder cells for which we have shown previously profen-induced apoptotic nuclear fragmentation (25). Treatment of the two prostate cancer cell lines (DU-145 and PC-3) with carprofen induced a dose-dependent (0–100  $\mu\text{mol/L}$ ) fragmentation of nuclei (Fig. 4). As negative controls, the MCF-7 and 3T3 cells that were not induced by carprofen to express p75<sup>NTR</sup> (Fig. 2A) did not undergo carprofen-dependent apoptotic nuclear fragmentation (Fig. 4). Expression of the  $\Delta\text{ICDp75}^{\text{NTR}}$  dominant-negative vector before carprofen treatment partially rescued nuclear fragmentation in the PC-3 and DU-145 prostate cells, whereas the MCF-7 and 3T3 negative control cells did not exhibit fragmented nuclei (Fig. 4).



**Figure 3.** PC-3 (A) and DU-145 (B) cell survival analysis by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay following 48 h treatment with 0  $\mu\text{mol/L}$  (cross-hatched), 50  $\mu\text{mol/L}$  (diagonal), or 100  $\mu\text{mol/L}$  (stippled) carprofen (Carp). Before treatment, cells were cotransfected with a ponasterone A-inducible ecdysone receptor plasmid pVgRxR and  $\Delta\text{ICDp75}^{\text{NTR}}$ . Following transfection, cells were incubated in serum-containing medium for 18 h and then incubated in 1  $\mu\text{mol/L}$  ponasterone A (P) for 24 h to drive expression of the dominant-negative gene products. Results are expressed relative to the control (0  $\mu\text{mol/L}$ ). \*,  $P < 0.001$ .

**Figure 4.** Detection of apoptotic nuclei (arrows) by Hoechst staining of PC-3 and DU-145 prostate cancer cells, MCF-7 breast cancer cells, and 3T3 fibroblasts following treatment with 0, 50, or 100  $\mu\text{mol/L}$  carprofen for 48 h. Before treatment, some cells were cotransfected with a ponasterone A-inducible ecdysone receptor plasmid pVgR $\times$ R and  $\Delta\text{ICDp75}^{\text{NTR}}$ . Following transfection, cells were incubated in serum-containing medium for 18 h and then incubated in 1  $\mu\text{mol/L}$  ponasterone A for 24 h to drive expression of the dominant-negative gene products and then treated with 100  $\mu\text{mol/L}$  carprofen for 48 h. Bar, 5  $\mu\text{m}$ .



#### Carprofen Induction of $p75^{\text{NTR}}$ Occurs via the p38 MAPK Pathway


An earlier study from our laboratory (24) implicated the aryl propionic acids, *R*-flurbiprofen and ibuprofen, in the induction of  $p75^{\text{NTR}}$  via the p38 MAPK pathway. Because carprofen, an aryl propionic acid, exhibits an order of magnitude greater potency (Fig. 2A) than *R*-flurbiprofen and ibuprofen for the induction of  $p75^{\text{NTR}}$  expression levels (23, 24), we examined the effect of siRNA knockdown of the p38 $\alpha$  MAPK isoform on  $p75^{\text{NTR}}$  levels following treatment with carprofen. We showed previously that p38 $\alpha$  MAPK is the predominant isoform expressed in PC-3 and DU-145 cells (24). Whereas treatment with carprofen induced  $p75^{\text{NTR}}$  expression levels, transfection of prostate cancer cells with p38 $\alpha$  siRNA before carprofen treatment prevented induction of  $p75^{\text{NTR}}$  relative to untransfected cells or cells transfected with nontargeting siRNA (Fig. 5A).

Because the p38 MAPK is activated by phosphorylation, we determined the phosphorylation status of p38 MAPK at several time points in PC-3 and DU-145 cells following treatment with carprofen. In both cell lines, carprofen treatment stimulated rapid phosphorylation of p38 MAPK as early as within 1 min treatment and subsequently led to the sustained activation of the p38 MAPK pathway that could be observed even 8 h following treatment of each cell line (Fig. 5B).

Because the MK2 kinase is downstream of p38 MAPK and was shown previously to be involved in profen induction of  $p75^{\text{NTR}}$  (24), we used a dominant-negative expression vector for MK2 to determine involvement in carprofen induction of  $p75^{\text{NTR}}$ . Treatment with carprofen alone induced expression of  $p75^{\text{NTR}}$  in both PC-3 and DU-145 cells, whereas transfection of dominant-negative MK2 before carprofen treatment decreased the induction of  $p75^{\text{NTR}}$  (Fig. 6).

#### Discussion

Carprofen is a propionic acid NSAID that induced  $p75^{\text{NTR}}$  levels in prostate cancer cell lines with an order of magnitude greater efficacy than the related propionic acid NSAIDs, *R*-flurbiprofen and ibuprofen (23). Concomitant with the superior efficacy of carprofen to induce levels of  $p75^{\text{NTR}}$  was its activity to inhibit cell survival via apoptosis. Our previous studies have shown a strong cause-and-effect relationship between induced levels of  $p75^{\text{NTR}}$  and induction of apoptosis in cancer cell lines (23, 25). When expression levels are induced,  $p75^{\text{NTR}}$  appears to be a robust marker of drug-induced apoptosis (23, 25). Because carprofen exhibited some degree of cell-specific induction of  $p75^{\text{NTR}}$ -associated apoptosis, we focused on the PC-3 and DU-145 prostate cancer cell lines, which were most responsive to carprofen treatment, and coincidentally are the




**Figure 5.** **A**, knockdown of p38 MAPK prevents induction of p75<sup>NTR</sup> by carprofen. PC-3 and DU-145 cells were transfected with nontargeting siRNA or siRNA for p38α for 72 h. Following transfection, cells were treated with 100 μmol/L carprofen (CAR) or DMSO vehicle control (CON) and the cell lysates were used for immunoblot analysis. A875 cell lysates were used as a positive control for p75<sup>NTR</sup> expression. β-Actin was used as the loading control. **B**, activation of the p38 MAPK pathway by carprofen. PC-3 and DU-145 cells were treated with 100 μmol/L carprofen for 0 min, 1 min, 5 min, 1 h, 4 h, or 8 h. Cell lysates were prepared for immunoblot analysis using antibodies to phosphorylated p38 MAPK (P-p38). Blots for phosphorylated p38 MAPK were stripped and reprobed for total p38 MAPK.

only two prostate cancer cell lines included in the NIH Developmental Therapeutics Program Anticancer Drug Discovery Program, due to their well-characterized aggressive phenotype. Hormone-responsive prostate cells were intentionally not included in these studies to maintain a focus on potential therapeutics of prostate tumor cells with phenotypes refractory to hormone ablation treatment consistent with poor prognosis. Using the prostate cancer cell lines, PC-3 and DU-145, most responsive to carprofen, we showed that a dominant-negative antagonist of p75<sup>NTR</sup> (ΔICDp75<sup>NTR</sup>) partially rescued carprofen-induced inhibition of cell survival, thereby confirming a cause-and-effect relationship between carprofen induction of p75<sup>NTR</sup> levels and p75<sup>NTR</sup> induction of apoptosis. Partial rather than complete rescue may be attributed to assay conditions or additional effects of carprofen independent of p75<sup>NTR</sup>.

In prostate cancer cell lines, reexpression of p75<sup>NTR</sup> induces modifications to several downstream signal transduction cascades leading to apoptosis. Initially, p75<sup>NTR</sup> expression down-regulates components of the nuclear factor-κB and c-Jun NH<sub>2</sub>-terminal kinase pathways preventing nuclear translocation of both these prosurvival transcriptional effectors (28). Expression of p75<sup>NTR</sup> also retards cell cycle progression through accumulation of cells in G<sub>1</sub> at the expense of S-phase cells (7, 26). Down-regulation of cyclin/cyclin-dependent kinase holoenzyme

components cyclin E, cyclin A, cyclin-dependent kinase 2, and cyclin-dependent kinase 6 contributes to hypophosphorylation of retinoblastoma along with elevated levels of p16<sup>INK4a</sup> in the p75<sup>NTR</sup>-induced cytostatic cells (26). Reexpression of p75<sup>NTR</sup> also induces elevated expression of the retinoic acid receptor β and retinoid X receptors α and β during partial redifferentiation of PC-3 cells that may also contribute to cytostasis (29). Evidence for p75<sup>NTR</sup>-dependent activation of extrinsic apoptosis in prostate cells has been limited to caspase-8 reductions in RIP, an adaptor protein that interacts with the ICD of p75<sup>NTR</sup> (28). Evidence for p75<sup>NTR</sup>-dependent activation of the intrinsic mitochondrial pathway includes an increase in proapoptotic effectors, Smac, Bax, Bak, and Bad, and conversely a decrease in the prosurvival effector, Bcl-xL (26), leading to a reduction in X-linked inhibitor of apoptosis protein and cleavage of caspase-9 and caspase-7 followed by poly(ADP-ribose) polymerase cleavage and nuclear fragmentation in PC-3 cells (26). Hence, reexpression of p75<sup>NTR</sup> appears to promote partial redifferentiation, cell cycle arrest, and apoptosis in prostate cancer cells, thereby providing a rationale for investigation of compounds that may be used for p75<sup>NTR</sup>-dependent therapeutics.

Prostate cancer cells evade the apoptotic effects of p75<sup>NTR</sup> expression by loss of p75<sup>NTR</sup> mRNA stability with concomitant suppression of p75<sup>NTR</sup> protein levels (6). Conversely, R-flurbiprofen and ibuprofen stabilize p75<sup>NTR</sup> mRNA with concomitant expression of p75<sup>NTR</sup> protein (24) and induction of apoptosis (23) through the p38 MAPK pathway (24). Indeed, abundant evidence has been reported for the involvement of p38 MAPK in apoptosis induced by a variety of agents such as the profen NSAIDs (23, 24), Fas ligation (30), and nerve growth factor withdrawal (31). The latter is significant because nerve growth factor ligation to the p75<sup>NTR</sup> acts as a survival signal in prostate cancer cells (26). Conversely, a relative absence of nerve growth factor, either by ligand withdrawal or by



**Figure 6.** PC-3 and DU-145 cells were transfected with dominant-negative MK2 (dnMK2), after which serum-containing medium was added for 24 h to allow expression of the dominant-negative gene product. Cells were treated with 100 μmol/L carprofen for 48 h. Cell lysates were collected for immunoblot analysis of p75<sup>NTR</sup> levels. β-Actin was used as the loading control.



up-regulation of p75<sup>NTR</sup> protein to levels that initially bind residual ligand and then to higher levels that result in unbound (no ligand) p75<sup>NTR</sup>, acts as a stimulus of apoptosis in prostate cancer cells (23, 26, 28). From a potential therapeutic perspective, this mechanism of p75<sup>NTR</sup>-dependent apoptosis has the appeal that agents, such as the profens, which elevate p75<sup>NTR</sup> levels, have the same effect as ligand withdrawal leading to apoptosis of cancer cells. Activation of the p38 MAPK signal transduction pathway by carprofen was rapid, within 1 min, suggesting that carprofen is interacting with a molecule highly proximal to p38 MAPK. The observation that p38 MAPK knockdown prevented carprofen induction of p75<sup>NTR</sup> levels confirms this pathway as a mechanism responsible for p75<sup>NTR</sup> regulation. We recently reported similar observations for *R*-flurbiprofen and ibuprofen activation of p38 MAPK up-regulation of p75<sup>NTR</sup>-dependent apoptosis in prostate cancer cells (24). In this pathway, MK2 directly binds to the p38 $\alpha$  isoform of MAPK during activation (32). Expression levels of MK2 are relatively robust in both PC-3 and DU-145 prostate cancer cells (24). Indeed, dominant-negative antagonism of MK2 prevented carprofen induction of p75<sup>NTR</sup> levels in prostate cancer cells. These observations suggest that carprofen initiates p75<sup>NTR</sup>-dependent apoptosis through a similar p38 MAPK signal transduction pathway to that of *R*-flurbiprofen and ibuprofen, albeit at an order of magnitude lower concentration of drug. Additional studies of this mechanism may lead to more potent compounds that induce p75<sup>NTR</sup>-dependent apoptosis of prostate cancer cells.

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Acknowledgments

We thank Prof. M. Gaestel for providing the dominant-negative MK2 construct.

## References

- Chao MV. The p75 neurotrophin receptor. *J Neurobiol* 1994;25:1373–85.
- Chapman BS. A region of the 75kD neurotrophin receptor homologous to the death domains of TNFR-1 and Fas. *FEBS Lett* 1995;374:216–20.
- Pflug BR, Dionne CA, Kaplan DR, et al. Expression of the Trk high affinity nerve growth factor receptor in the human prostate. *Endocrinology* 1995;136:262–8.
- Perez M, Regan T, Pflug B, et al. Loss of the low affinity nerve growth factor receptor during malignant transformation of the human prostate. *Prostate* 1997;30:274–9.
- Pflug BR, Onoda M, Lynch JH, et al. Reduced expression of the low affinity nerve growth factor receptor in benign and malignant human prostate tissue and loss of expression in four human metastatic prostate tumor cell lines. *Cancer Res* 1992;52:5403–6.
- Krygiel S, Djakiew D. Molecular characterization of p75<sup>NTR</sup> loss of the expression in human prostate tumor cells. *Mol Carcinogenesis* 2001;31:46–55.
- Krygiel S, Djakiew D. The neurotrophin receptor p75<sup>NTR</sup> is a tumor suppressor in human prostate cancer. *Anticancer Res* 2001;21:3749–55.
- Krygiel S, Djakiew D. Neurotrophin receptor p75<sup>NTR</sup> suppresses growth and nerve growth factor-mediated metastasis of human prostate cancer cells. *Int J Cancer* 2002;98:1–7.
- Takeito MM. Cyclooxygenase-2 inhibitors in tumorigenesis. Part II. *J Natl Cancer Inst* 1998;90:1609–14.
- Vainio H. Is COX-2 inhibition a panacea for cancer prevention? *Int J Cancer* 2001;94:613–8.
- Nelson JE, Harris RE. Inverse association of prostate cancer and non-steroidal anti-inflammatory drugs (NSAIDs): results of a case-control study. *Oncol Rep* 2000;7:169–73.
- Norrish AE, Jackson RT, McRae CU. Non-steroidal anti-inflammatory drugs and prostate cancer progression. *Int J Cancer* 1998;77:511–4.
- Roberts RO, Jacobson DJ, Gorman CJ, et al. A population-based study of daily nonsteroidal anti-inflammatory drug use and prostate cancer. *Mayo Clinic Proc* 2002;77:219–23.
- Drago JR, Murray C. Control of metastases in the Nb rat prostatic adenocarcinoma model. *J Androl* 1984;5:265–8.
- Liu XH, Kirschenbaum A, Yao S, et al. Inhibition of cyclooxygenase-2 suppresses angiogenesis and growth of prostate cancer *in vivo*. *J Urol* 2000;164:820–4.
- Pollard M, Chang CF. Investigations on prostatic adenocarcinomas in rats. *Oncology* 1977;34:129–33.
- Vane JR. Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nat New Biol* 1971;231:232–5.
- Sheng H, Shao J, Morrow JD, et al. Modulation of apoptosis and Bcl-2 expression by prostaglandin E<sub>2</sub> in human colon cancer cells. *Cancer Res* 1998;58:362–6.
- Masferrer JL, Leahy KM, Koki AT, et al. Antiangiogenic and antitumor activities of cyclooxygenase-2 inhibitors. *Cancer Res* 2000;60:1306–11.
- Zhang X, Morham SG, Langenbach R, et al. Malignant transformation and antineoplastic actions of nonsteroidal antiinflammatory drugs (NSAIDs) on cyclooxygenase-null embryo fibroblasts. *J Exp Med* 1999;190:451–9.
- Wechter WJ, Kantoci D, Murray ED, Jr., et al. *R*-flurbiprofen chemoprevention and treatment of intestinal adenomas in the APC(Min)/+ mouse model: implications for prophylaxis and treatment of colon cancer. *Cancer Res* 1997;57:4316–24.
- Lim JT, Piazza, GA, Han EK, et al. Sulindac derivatives inhibit growth and induce apoptosis in human prostate cancer cell lines. *Biochem Pharm* 1999;58:1097–107.
- Quann E, Khwaja F, Zavitz KH, et al. The aryl propionic acid *R*-flurbiprofen selectively induces p75<sup>NTR</sup>-dependent decreased survival of prostate tumor cells. *Cancer Res* 2007;67:3254–62.
- Quann E, Khwaja F, Djakiew D. The p38 MAPK pathway mediates aryl propionic acid-induced messenger RNA stability of p75<sup>NTR</sup> in prostate cancer cells. *Cancer Res* 2007;67:11402–10.
- Khwaja F, Allen J, Lynch J, et al. Ibuprofen inhibits survival of bladder cancer cells by induced expression of the p75<sup>NTR</sup> tumor suppressor protein. *Cancer Res* 2004;64:6207–13.
- Khwaja F, Tabassum A, Allen J, et al. The p75(NTR) tumor suppressor induces cell cycle arrest facilitating caspase mediated apoptosis in prostate tumor cells. *Biochem Biophys Res Commun* 2006;341:1184–92.
- Engel KH, Schultz F, Martin F, et al. Constitutive activation of mitogen-activated protein kinase-activated protein kinase 2 by mutation of phosphorylation sites and an A-helix motif. *J Biol Chem* 1995;270:27213–21.
- Allen J, Khwaja F, Byers S, et al. The p75<sup>NTR</sup> mediates a bifurcated signal transduction cascade through the NF $\kappa$ B and JNK pathways to inhibit cell survival. *Exp Cell Res* 2005;304:69–80.
- Nalbandian A, Pang AL, Rennert OM, et al. A novel function of differentiation revealed by cDNA microarray profiling of p75<sup>NTR</sup>-regulated gene expression. *Differentiation* 2005;73:385–96.
- Juo P, Kuo CJ, Reynolds, et al. Fas activation of the p38 mitogen-activated protein kinase signaling pathway requires ICE/CED-3 family proteases. *Mol Cell Biol* 1997;17:24–35.
- Xia Z, Dickens M, Raingeaud J, et al. Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 1995;270:1326–31.
- Gaestel M. MAPKAP kinases, MKs, two's company, three a crowd. *Mol Cell Biol* 2006;26:120–30.

# The p38 MAPK Pathway Mediates Aryl Propionic Acid–Induced Messenger RNA Stability of p75<sup>NTR</sup> in Prostate Cancer Cells

Emily J. Quann,<sup>1</sup> Fatima Khwaja,<sup>1</sup> and Daniel Djakiew<sup>1,2</sup>

<sup>1</sup>Department of Biochemistry and Molecular & Cellular Biology and <sup>2</sup>Vincent T. Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, District of Columbia

## Abstract

The p75<sup>NTR</sup> acts as a tumor suppressor in the prostate, but its expression is lost as prostate cancer progresses and is minimal in established prostate cancer cell lines such as PC-3, DU-145, and LNCaP. Previously, we showed that treatment with R-flurbiprofen or ibuprofen induced p75<sup>NTR</sup> expression in PC-3 and DU-145 cells leading to p75<sup>NTR</sup>-mediated decreased survival. Here, we investigate the mechanism by which these drugs induce p75<sup>NTR</sup> expression. We show that the observed increase in p75<sup>NTR</sup> protein due to R-flurbiprofen and ibuprofen treatment was accompanied by an increase in p75<sup>NTR</sup> mRNA, and this increase in mRNA was the result of increased mRNA stability and not by an up-regulation of transcription. In addition, we show that treatment with R-flurbiprofen or ibuprofen led to sustained activation of the p38 mitogen-activated protein kinase (MAPK) pathway. Furthermore, inhibition of the p38 MAPK pathway with the p38 MAPK-specific inhibitor SB202190 or by small interfering RNA (siRNA) knockdown of p38 MAPK protein prevented induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen. We also observed that siRNA knockdown of MAPK-activated protein kinase (MK)-2 and MK3, the kinases downstream of p38 MAPK that are responsible for the mRNA stabilizing effects of the p38 MAPK pathway, also prevented an induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen. Finally, we identify the RNA stabilizing protein HuR and the posttranscriptional regulator eukaryotic translation initiation factor 4E as two possible mechanisms by which the p38 MAPK pathway may increase p75<sup>NTR</sup> expression. Collectively, the data suggest that R-flurbiprofen and ibuprofen induce p75<sup>NTR</sup> expression by increased mRNA stability that is mediated through the p38 MAPK pathway. [Cancer Res 2007;67(23):11402–10]

## Introduction

The p75<sup>NTR</sup> (neurotrophin receptor) is a member of the tumor necrosis factor (TNF) receptor superfamily that share a conserved intracellular death domain capable of inducing apoptosis and suppressing growth (1, 2). It binds neurotrophin ligands with similar affinity; however, unlike other members of the TNF receptor superfamily, it is able to induce apoptosis and suppress growth in the unbound state (3–8). p75<sup>NTR</sup> has been shown to act as a tumor suppressor in the prostate (5, 7, 9). However, whereas normal

prostate epithelial cells express high levels of p75<sup>NTR</sup>, expression is lost as prostate cancer progresses (10). In addition, although the gene remains intact, expression is very low in metastatic prostate cancer cell lines PC-3, DU-145, and LNCaP (10). Exogenous reexpression of p75<sup>NTR</sup> in PC-3 cells led to increased apoptosis, reduced proliferation, and a decreased ability of these cells to form tumors in mice, thus indicating potential for treatments that result in the reexpression of p75<sup>NTR</sup> in prostate tumor cells (5, 7–9, 11).

Nonsteroidal anti-inflammatory drugs are typically used to relieve inflammation by inhibiting cyclooxygenase (COX) activity; however, many of these drugs seem to possess anticancer activity that is independent of their COX inhibitory activity (12). Flurbiprofen and ibuprofen are members of the aryl propionic acid class of nonsteroidal anti-inflammatory drugs, also referred to as profens, and have shown anticancer activity in the prostate. For example, treatment with the enantiomer R-flurbiprofen, which lacks COX inhibitory activity, was able to slow progression of prostate cancer in the transgenic adenocarcinoma of the mouse prostate mouse (13). In addition, long-term ibuprofen use has been associated with decreased prostate cancer risk, and treatment of prostate cancer cells with ibuprofen resulted in decreased survival (14–16). Recently, we showed that treatment of PC-3 and DU-145 prostate cancer cells with R-flurbiprofen or ibuprofen resulted in a strong induction of p75<sup>NTR</sup>, which led to p75<sup>NTR</sup>-mediated apoptosis and decreased survival (17). The results of this study were significant because they showed that p75<sup>NTR</sup> expression is inducible, and that drugs which induce p75<sup>NTR</sup> in prostate cancer cells have therapeutic potential.

The p38 mitogen-activated protein kinase (MAPK) pathway is activated in response to cell stresses and external stimuli such as heat, UV light, osmotic shock, and inflammatory cytokines (18). It mediates various cellular processes including apoptosis, senescence, inflammation, and tumorigenesis through decreased p38 MAPK activity, which has been associated with tumor progression (18–20). In addition, p38 MAPK has been strongly implicated as a regulator of mRNA stability (18, 21, 22). p38 MAPK is activated through phosphorylation by upstream kinases MAPK kinase kinase (MKK)-3 and MKK6 and, on activation, phosphorylates a number of downstream targets (18). Among these targets are the kinases MAPK-activated protein kinase (MK)-2 and MK3, which have been shown to be responsible for mediating the effects of the p38 MAPK pathway on mRNA stability (21, 23).

In this study, we investigated the mechanism by which the profens, R-flurbiprofen and ibuprofen, induce p75<sup>NTR</sup> expression in PC-3, DU-145, and LNCaP prostate cancer cells. We showed that induction of p75<sup>NTR</sup> protein expression corresponded to increased p75<sup>NTR</sup> mRNA levels that were due to increased mRNA stability and not to an up-regulation of transcription. In addition, we showed that R-flurbiprofen and ibuprofen treatment led to sustained activation of the p38 MAPK pathway and that this pathway was necessary for induction of p75<sup>NTR</sup> expression.

**Note:** Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org/>).

**Requests for reprints:** Daniel Djakiew, Department of Biochemistry and Molecular & Cellular Biology, Georgetown University Medical Center, 3900 Reservoir Road Northwest, Washington, DC 20057-1436. Phone: 202-687-1203; Fax: 202-687-1823; E-mail: djakiewd@georgetown.edu.

©2007 American Association for Cancer Research.  
doi:10.1158/0008-5472.CAN-07-1792

## Materials and Methods

**Cell lines, culture conditions, and reagents.** PC-3 and DU-145 cells, the only two prostate tumor cell lines included in the NIH Developmental Therapeutics Program anticancer drug discovery program, and the androgen receptor-responsive LNCaP human prostate cancer cells were purchased from the tissue culture core facility of the Georgetown University Lombardi Comprehensive Cancer Center and maintained as previously described (17).

Ibuprofen (Sigma Chemical Co.) and R-flurbiprofen (Myriad Pharmaceuticals, Inc.) were dissolved in DMSO and used at final concentrations of 0, 0.25, 0.5, 1.0, or 2.0 mmol/L. Actinomycin D (Tocris Bioscience) and SB202190 (Calbiochem) were dissolved in DMSO and used at final concentrations of 5 µg/mL and 20 µmol/L, respectively.

**Reverse transcription-PCR.** RNA was isolated using TRIzol reagent (Invitrogen) according to the manufacturer's protocol. Reverse transcription-PCR (RT-PCR) was done with the SuperScript III One-Step RT-PCR System with Platinum Taq DNA Polymerase (Invitrogen) using equal amounts of RNA according to the following program: 47°C for 30 min; 94°C for 2 min; 30 cycles of 94°C for 1 min, 60°C for 1 min, and 72°C for 1 min; and 72°C for 5 min. Primers for p75<sup>NTR</sup> and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) were described previously (17).

**Immunoblot analysis.** Lysates were prepared as described previously (11). Cytoplasmic and nuclear fractions were prepared with a nuclear extract kit (Active Motif). SDS-PAGE and immunoblot analysis were done as described previously (17). The following primary antibodies were used: mouse monoclonal anti-p75<sup>NTR</sup> (Millipore; 1:2,000); rabbit polyclonal antibodies to phosphorylated p38 MAPK, p38 MAPK, p38α, phosphorylated MK2, MK2, phosphorylated eukaryotic translation initiation factor 4E (eIF4E), and eIF4E (Cell Signaling Technology; 1:1,000); rabbit polyclonal anti-HuR (Millipore; 1:1,000); mouse monoclonal anti-p38β (Zymed Laboratories) and anti-MK3 (Affinity BioReagents; 2 µg/mL); mouse monoclonal anti-Hu protein (Abcam, Inc.; 1:1,000); mouse monoclonal anti-α-tubulin and rabbit polyclonal anti-lamin B1 (Abcam; 1:3,000); and mouse monoclonal anti-β-actin (Sigma; 1:5,000). Goat anti-mouse and goat anti-rabbit horseradish peroxidase-conjugated secondary antibodies (Bio-Rad Laboratories) were used at 1:3,000. Membranes used for detection of phosphorylated p38 MAPK, MK2, or eIF4E were subsequently stripped and reprobed for total p38 MAPK, MK2, or eIF4E, respectively.

**Small interfering RNA transfection.** Cells were transfected for 72 h with nontargeting small interfering RNA (siRNA) or siRNA specific for p38α (J-003512-20), p38β (J-003972-11), MK2 (J-003516-11), MK3 (J-005014-07), or HuR (J-003773-08; Dharmacon RNA Technologies) at final concentrations of 100 nmol/L according to the manufacturer's protocol. Transfection reagent DharmaFECT 1 was used for DU-145 cells and DharmaFECT 2 was used for PC-3 and LNCaP cells (Dharmacon).

**Luciferase assay.** PC-3, DU-145, and A875 cells were cotransfected with either the pGL3-2.1 kb p75<sup>NTR</sup> promoter-luciferase construct (gift from Emil Bogenmann, Children's Hospital, Los Angeles, CA) or empty vector and the *Renilla* luciferase reporter vector pRL-TK (Promega) using GeneJammer transfection reagent (Stratagene) for 48 h. Following transfection, PC-3 and DU-145 cells were treated with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control for 24 h, whereas A875 cells were treated with isotonic medium (320 mosm/L) or hypoosmotic medium (160 mosm/L) prepared by a 1:1 dilution of complete medium with water. Luciferase activity was measured using the Dual-Luciferase Reporter Assay (Promega) and normalized to the empty vector and *Renilla* luciferase activity.

**Immunoprecipitation RT-PCR.** Equal numbers of treated cells were collected, washed in cold PBS, and incubated on ice for 30 min in 1 mL of lysis buffer (1% Igepal Ca-630, 0.5% sodium deoxycholate, 0.1% SDS in PBS) containing 10 µg/mL Protease Inhibitor Cocktail (Sigma) and 100 units of RNaseOUT (Invitrogen) per sample. Samples were then centrifuged at 10,000 × g for 15 min at 4°C. Ten micrograms of HuR antibody (Millipore) were added to the supernatant of each sample, with the exception of the negative controls, and samples were rocked at 4°C for 30 min. Fifty microliters of protein A/G Plus-agarose beads (Santa Cruz Biotechnologies)

were added to each sample, and samples were rocked at 4°C for 30 min. Samples were centrifuged at 2,000 × g for 2 min at 4°C, the supernatant was discarded, and the beads were washed eight times with 1 mL of lysis buffer. RNA was isolated from the beads using TRIzol reagent (Invitrogen) according to the manufacturer's protocol. RT-PCR was done as described above for determination of p75<sup>NTR</sup> mRNA level using equal volumes of RNA and 32 cycles of PCR.


## Results

### R-Flurbiprofen and ibuprofen increase p75<sup>NTR</sup> mRNA level.

To determine the mechanism by which the profens, R-flurbiprofen and ibuprofen, induce p75<sup>NTR</sup> expression in PC-3 and DU-145 human prostate cancer cell lines, we first compared the levels of p75<sup>NTR</sup> protein and mRNA following treatment with various concentrations of each profen (Fig. 1A). There was a strong correlation between p75<sup>NTR</sup> protein and mRNA, indicating that increased mRNA levels are responsible for the induction of p75<sup>NTR</sup> protein by R-flurbiprofen and ibuprofen. A strong increase in p75<sup>NTR</sup> protein and mRNA levels was observed in PC-3 cells following treatment with 0.5 mmol/L R-flurbiprofen and 1.0 mmol/L ibuprofen, and in DU-145 cells with 1.0 mmol/L R-flurbiprofen and 2.0 mmol/L ibuprofen. Therefore, in subsequent experiments, we used 2.0 mmol/L R-flurbiprofen and ibuprofen because at this concentration, each of these profens induced high levels of p75<sup>NTR</sup> in both cell types. The level of p75<sup>NTR</sup> mRNA was also determined at various time points following profen treatment, and an induction was first observed within 2 to 4 h of treatment (Fig. 1B). We also compared levels of p75<sup>NTR</sup> protein and mRNA in the androgen-responsive LNCaP cell line following treatment with several concentrations of ibuprofen, and again observed a strong correlation between p75<sup>NTR</sup> protein and mRNA (Supplementary Fig. S1A). In addition, we observed a similar time course for induction of p75<sup>NTR</sup> mRNA in LNCaP cells treated with ibuprofen (Supplementary Fig. S1B).

### R-Flurbiprofen and ibuprofen increase p75<sup>NTR</sup> mRNA stability.

Because the amount of transcript in a cell is a function of the rate of transcription and the rate of degradation, the observed increase in p75<sup>NTR</sup> mRNA in PC-3 and DU-145 cells following R-flurbiprofen and ibuprofen treatment must be due to either an up-regulation of transcriptional activity or a decrease in the rate of degradation of the p75<sup>NTR</sup> transcript, or both. Thus, we examined transcriptional activity and mRNA stability following treatment with profens. PC-3 and DU-145 cells were transfected with a p75<sup>NTR</sup> promoter-luciferase construct followed by treatment with R-flurbiprofen or ibuprofen, and promoter activity was compared with DMSO control-treated cells (Fig. 2A). Treatment with profens did not result in a significant change in p75<sup>NTR</sup> promoter activity, suggesting that the increase in mRNA was not due to transcriptional up-regulation of the p75<sup>NTR</sup> gene. Previously, p75<sup>NTR</sup> promoter activity was shown to be up-regulated in the A875 melanoma cell line on exposure to hypoosmotic medium (24). Therefore, to show that the p75<sup>NTR</sup> promoter-luciferase construct was functional, we transfected A875 cells with the construct and compared luciferase activity in cells exposed to isotonic versus hypoosmotic medium. Consistent with previously published data, treatment with hypoosmotic medium resulted in 17-fold increased p75<sup>NTR</sup> promoter activity (Fig. 2A; ref. 24). Therefore, increased levels of p75<sup>NTR</sup> mRNA and protein in PC-3 and DU-145 cells treated with R-flurbiprofen or ibuprofen were not a consequence of changed transcriptional activity.



**Figure 1.** A, p75<sup>NTR</sup> mRNA and protein levels following R-flurbiprofen and ibuprofen treatment. PC-3 and DU-145 cells were treated with 0 to 2 mmol/L R-flurbiprofen or ibuprofen for 24 h. Cell lysates were collected and equal amounts of protein were subjected to SDS-PAGE and immunoblot analysis with an antibody to p75<sup>NTR</sup> or β-actin (β-act) for the loading control. Alternatively, RNA was isolated from PC-3 and DU-145 cells after 24 h of R-flurbiprofen or ibuprofen treatment, and p75<sup>NTR</sup> and GAPDH mRNA levels were determined by RT-PCR. B, time course induction of p75<sup>NTR</sup> mRNA levels. PC-3 and DU-145 cells were treated with 2 mmol/L R-flurbiprofen or ibuprofen for 0, 2, 4, 8, or 24 h. RNA was isolated at each time point and levels of p75<sup>NTR</sup> and GAPDH were determined by RT-PCR.

The stability of the p75<sup>NTR</sup> transcript was determined with the transcriptional inhibitor actinomycin D. PC-3 and DU-145 cells were treated with R-flurbiprofen, ibuprofen, or DMSO for 8 h followed by the addition of actinomycin D. p75<sup>NTR</sup> mRNA level was determined by RT-PCR at several time points following actinomycin D addition (Fig. 2B and C). The majority of p75<sup>NTR</sup> mRNA from DMSO control-treated PC-3 and DU-145 cells was degraded within 2 and 4 h, respectively. In contrast, p75<sup>NTR</sup> mRNA was still present at high levels in profen-treated cells 12 h following the addition of actinomycin D. Therefore, profen treatment strongly enhanced p75<sup>NTR</sup> mRNA stability but did not influence transcription, suggesting that increased mRNA stability is responsible for the induction of p75<sup>NTR</sup> expression by R-flurbiprofen and ibuprofen (Fig. 2). We also observed an increase in p75<sup>NTR</sup> mRNA stability in LNCaP cells treated with ibuprofen relative to DMSO control-treated cells (Supplementary Fig. S1C and D).

**R-Flurbiprofen and ibuprofen activate the p38 MAPK pathway.** The p38 MAPK pathway has been strongly implicated in the stabilization of several mRNAs through the downstream kinases MK2 and MK3, which are activated upon phosphorylation by p38 MAPK (21, 23). p38 MAPK is also activated by phosphorylation. Therefore, we determined the phosphorylation status of p38 MAPK and MK2 at several time points in PC-3 and DU-145 cells following treatment with R-flurbiprofen or ibuprofen and in LNCaP cells following treatment with ibuprofen (Fig. 3A and Supplementary Fig. S2A). In all cases, profen treatment led to sustained activa-

tion of the p38 MAPK pathway, which could be observed even 8 h after treatment of each cell line.

**The p38 MAPK pathway is involved in the induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen.** Increased p75<sup>NTR</sup> mRNA stability seems to be the mechanism by which R-flurbiprofen and ibuprofen induce p75<sup>NTR</sup> expression, and these drugs activate the p38 MAPK pathway, which is known to be involved in mRNA stabilization. Pretreatment of PC-3 and DU-145 cells with the p38 MAPK inhibitor SB202190 followed by treatment with R-flurbiprofen or ibuprofen prevented an induction of p75<sup>NTR</sup>, indicating that p38 MAPK is involved in profen-mediated induction of p75<sup>NTR</sup> (Fig. 3B). To further confirm the role of the p38 MAPK pathway in profen-mediated induction of p75<sup>NTR</sup>, we also used p38 MAPK siRNA to specifically knock down p38 MAPK protein. There are two ubiquitously expressed isoforms of p38 MAPK, p38α and p38β, which are capable of phosphorylating MK2 and MK3 (18, 23, 25). PC-3 and DU-145 cells were transfected with nontargeting siRNA, p38α siRNA, p38β siRNA, or siRNAs for p38α and p38β together. The efficacy and specificity of the siRNAs were determined by Western blot (Fig. 4A). Transfection of cells with p38α siRNA or p38α and p38β siRNAs together before profen treatment prevented an induction of p75<sup>NTR</sup> relative to untransfected cells or cells transfected with nontargeting siRNA (Fig. 4B). Transfection with p38β siRNA alone was less effective in preventing induction of p75<sup>NTR</sup> than transfection with p38α siRNA. Because the p38β isoform seems to be expressed at much lower

levels than p38 $\alpha$ , it is not surprising that knockdown of p38 $\beta$  alone only modestly affected induction of p75<sup>NTR</sup> because p38 $\alpha$  was still present. Expression of p38 $\alpha$  was also much greater than that of p38 $\beta$  in LNCaP cells (not shown), and consistently, induction of p75<sup>NTR</sup> by ibuprofen was prevented in LNCaP cells transfected with p38 $\alpha$  siRNA (Supplementary Fig. S2C).


Because the MK2 and MK3 kinases are downstream of p38 MAPK and have both been shown to mediate mRNA stabilization, we used siRNA for MK2 and MK3 to determine if they are involved in the induction of p75<sup>NTR</sup> by R-flurbiprofen or ibuprofen. PC-3 and DU-145 cells were transfected with nontargeting siRNA, MK2 siRNA, MK3 siRNA, or siRNAs for MK2 and MK3 in combination. The efficacy and specificity of these siRNAs were tested in PC-3 and DU-145 cells by Western blot (Fig. 5A). Transfection with MK2 siRNA or MK3 siRNA separately before profen treatment resulted in decreased induction of p75<sup>NTR</sup> (Fig. 5B). However, transfection with MK2 siRNA and MK3 siRNA together, resulting in the simultaneous knockdown of both proteins, was more effective in preventing induction of p75<sup>NTR</sup> by R-flurbiprofen or ibuprofen than

knockdown of MK2 or MK3 separately (Fig. 5B). This is consistent with previous observations that MK2 and MK3 are activated in parallel by p38 MAPK, have similar substrates, and are both strongly expressed in PC-3 and DU-145 cells (Fig. 5A; refs. 23, 25). Likewise, in LNCaP cells, simultaneous siRNA knockdown of MK2 and MK3 was successful in preventing induction of p75<sup>NTR</sup> expression by ibuprofen (Supplementary Fig. S2C).

#### p75<sup>NTR</sup> mRNA is a target of the RNA stabilizing protein HuR.

The HuR is an RNA binding protein that enhances the stability of target mRNAs (26, 27). It has been shown that activation of the p38 MAPK pathway results in the translocation of HuR from the nucleus to the cytoplasm where it binds to the 3'-untranslated regions (UTR) of various mRNAs containing a core AUUUA sequence (20, 28–31). The p75<sup>NTR</sup> transcript contains two AUUUA sites in the 3'-UTR, suggesting HuR binding as a possible mechanism by which R-flurbiprofen and ibuprofen may stabilize p75<sup>NTR</sup> mRNA. R-Flurbiprofen- and ibuprofen-treated cells exhibited modestly increased levels of cytoplasmic HuR (Fig. 6A). In addition, siRNA knockdown of HuR protein before profen treatment of PC-3 and

**Figure 2.** A, p75<sup>NTR</sup> promoter activity following R-flurbiprofen and ibuprofen treatment. PC-3 and DU-145 cells were transfected for 48 h with a p75<sup>NTR</sup> promoter-luciferase construct and then treated with 2 mmol/L R-flurbiprofen (FLU), 2 mmol/L ibuprofen (IBU), or DMSO vehicle control (C) for 24 h. Relative luciferase activity was determined after normalization to *Renilla* and graphed as fold induction relative to DMSO control-treated cells. As a positive control, A875 cells were transfected for 48 h with the p75<sup>NTR</sup> promoter-luciferase construct and then exposed to isotonic (Iso) or hypoosmotic (Hypo) medium for 24 h. Fold induction of luciferase activity was determined for hypoosmotic treated cells relative to cells in isotonic medium. Bars, SE. \*\*\*,  $P < 0.0001$  (ANOVA). B and C, increase in p75<sup>NTR</sup> mRNA stability following R-flurbiprofen or ibuprofen treatment. PC-3 and DU-145 cells were treated with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control for 8 h followed by the addition of 5  $\mu$ g/mL actinomycin D (ActD). RNA was isolated at 0, 2, 4, 8, and 12 h following the addition of actinomycin D, and p75<sup>NTR</sup> mRNA levels were determined by RT-PCR. The intensity of the bands for p75<sup>NTR</sup> was analyzed by densitometry and graphed as a percentage of the 0-h time point.






DU-145 cells partially prevented induction of  $p75^{\text{NTR}}$  relative to untransfected or nontargeting siRNA-transfected cells, suggesting that HuR is involved in the induction of  $p75^{\text{NTR}}$  by R-flurbiprofen and ibuprofen (Fig. 6B). Similarly, transfection of LNCaP cells with HuR siRNA before ibuprofen treatment was effective in preventing increased  $p75^{\text{NTR}}$  expression (Supplementary Fig. S2C). HuR belongs to the embryonic lethal abnormal vision family of proteins, which also includes HuB, HuC, and HuD, all RNA stabilizing proteins that are typically expressed in neurons (32). Whereas HuR expression was detected in PC-3 and DU-145 cells (Fig. 6A), expression of the other family members could not be detected (not shown), indicating that they most likely do not participate in profen-mediated induction of  $p75^{\text{NTR}}$ . A direct interaction between the HuR protein and the  $p75^{\text{NTR}}$  transcript was shown by an immunoprecipitation RT-PCR assay. Lysates of profen-treated PC-3 and DU-145 cells were immunoprecipitated with, or without, an HuR antibody and RNA was isolated from the immunoprecipitated beads, followed by RT-PCR for  $p75^{\text{NTR}}$  mRNA. PC-3 and DU-145 cells treated with R-flurbiprofen or ibuprofen showed a direct interaction between HuR and  $p75^{\text{NTR}}$  mRNA, whereas DMSO control-treated cells did not (Fig. 6C). In addition,  $p75^{\text{NTR}}$  mRNA was not detected in samples from cells treated with ibuprofen but not incubated with the HuR antibody, indicating that the  $p75^{\text{NTR}}$  mRNA detected in the treated samples was not nonspecific (Fig. 6C).

**R-Flurbiprofen and ibuprofen increase phosphorylation of eIF4E.** Although HuR seems to bind to the  $p75^{\text{NTR}}$  transcript upon profen treatment, the inhibition of  $p75^{\text{NTR}}$  induction in the

presence of a substantial HuR knockdown was modest. This result suggests that there are additional mechanisms involved in the induction of  $p75^{\text{NTR}}$  by the p38 MAPK pathway. eIF4E is a downstream, indirect target of p38 MAPK and is involved in several mechanisms of posttranscriptional regulation including translation initiation through 5'-cap binding, mRNA nuclear export, and mRNA stability (33, 34). We examined eIF4E phosphorylation following profen treatment in untransfected, nontargeting siRNA-transfected, and p38 $\alpha$  siRNA-transfected PC-3 and DU-145 cells. R-Flurbiprofen and ibuprofen treatment resulted in increased phosphorylated eIF4E levels in both cell types, and the increase was inhibited to varying degrees in the p38 $\alpha$  siRNA-transfected cells (Fig. 6D). These results identify eIF4E phosphorylation as another mechanism through which the p38 MAPK pathway may regulate  $p75^{\text{NTR}}$  expression.


## Discussion

The  $p75^{\text{NTR}}$  exhibits tumor suppressor activity in prostate cancer cells, which is mediated through an intracellular death domain capable of inducing apoptosis and suppressing growth (2, 5, 7, 9). However,  $p75^{\text{NTR}}$  expression is progressively lost in organ confined prostate cancer and is minimal in established metastatic prostate cancer cell lines (10). Significantly, our recent observations that the profens, R-flurbiprofen and ibuprofen, induced reexpression of  $p75^{\text{NTR}}$  protein levels causal of  $p75^{\text{NTR}}$ -dependent decreased survival of prostate cancer cells has stimulated an investigation



**Figure 3.** A, activation of the p38 MAPK pathway by R-flurbiprofen and ibuprofen. PC-3 and DU-145 cells were treated with 2 mmol/L R-flurbiprofen or 2 mmol/L ibuprofen for 0, 5 min, 1 h, 4 h, or 8 h. Cell lysates were collected and equal amounts of protein were subjected to SDS-PAGE and immunoblot analysis with antibodies to phosphorylated p38 MAPK (p-p38) or phosphorylated MK2 (p-MK2). Blots for phosphorylated p38 MAPK and phosphorylated MK2 were stripped and reprobed for total p38 MAPK and total MK2, respectively. β-actin was used as the loading control. B, the p38 MAPK inhibitor SB202190 prevents induction of  $p75^{\text{NTR}}$  by R-flurbiprofen and ibuprofen. PC-3 and DU-145 cells were pretreated with or without 20 μmol/L SB202190 (SB) for 1 h and then treated with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control (CON) for 24 h. Cell lysates were collected and equal amounts of protein were subjected to SDS-PAGE and immunoblot analysis with an antibody to  $p75^{\text{NTR}}$  or β-actin for the loading control.


**Figure 4.** A, validation assay for siRNA knockdown of p38 MAPK. PC-3 and DU-145 cells were transfected for 72 h with nontargeting siRNA or siRNA targeting p38 $\alpha$ , p38 $\beta$ , or siRNAs for p38 $\alpha$  and p38 $\beta$  together. Cell lysates were collected and efficacy of siRNA knockdown was determined by SDS-PAGE and immunoblot analysis with antibodies to p38 $\alpha$  and p38 $\beta$ .  $\beta$ -actin was used as the loading control. B, knockdown of p38 MAPK prevents induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen. PC-3 and DU-145 cells were transfected with nontargeting siRNA, siRNA for p38 $\alpha$ , siRNA for p38 $\beta$ , or siRNAs for p38 $\alpha$  and p38 $\beta$  together for 72 h. Following transfection, cells were treated with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control for 24 h. Cell lysates were collected and expression of p75<sup>NTR</sup> was determined by SDS-PAGE and immunoblot analysis.  $\beta$ -actin was used as the loading control.




of the associated signal transduction cascade that mediates profen effects on prostate cancer cells (17). Increased p75<sup>NTR</sup> protein levels in R-flurbiprofen- or ibuprofen-treated cells were accompanied by increased mRNA levels. This indicated that profen treatment

influenced either the rate of mRNA degradation or the rate at which the gene was transcribed. Profen-treated cells showed no significant change in luciferase activity of the p75<sup>NTR</sup> promoter construct relative to untreated cells. However, the transcriptional

**Figure 5.** A, validation assay for siRNA knockdown of MK2 and MK3. PC-3 and DU-145 cells were transfected for 72 h with nontargeting siRNA, siRNA for MK2, siRNA for MK3, or siRNAs for MK2 and MK3 together. Cell lysates were collected and efficacy of siRNA knockdown was determined by SDS-PAGE and immunoblot analysis with antibodies to MK2 and MK3.  $\beta$ -actin was used as the loading control. B, knockdown of MK2 and MK3 prevents induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen. PC-3 and DU-145 cells were transfected with nontargeting siRNA, siRNA for MK2, siRNA for MK3, or siRNAs for MK2 and MK3 together for 72 h. Following transfection, cells were treated with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control for 24 h. Cell lysates were collected and expression of p75<sup>NTR</sup> was determined by SDS-PAGE and immunoblot analysis.  $\beta$ -actin was used as the loading control.





**Figure 6.** A, translocation of HuR from the nucleus to the cytoplasm following treatment with R-flurbiprofen or ibuprofen. PC-3 and DU-145 cells were treated for 2 h with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control. Nuclear (N) and cytoplasmic (C) fractions were subjected to SDS-PAGE and immunoblot analysis with antibodies to HuR, lamin B1, and  $\alpha$ -tubulin. B, knockdown of HuR prevents induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen. PC-3 and DU-145 cells were transfected for 72 h with nontargeting siRNA or siRNA targeting HuR. For the test for siRNA knockdown, cell lysates were collected and efficacy of siRNA knockdown was determined by SDS-PAGE and immunoblot analysis with an antibody to HuR or  $\beta$ -actin for the loading control. For examination of p75<sup>NTR</sup> expression, cells were treated with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control for 24 h following siRNA transfection. Cell lysates were collected and expression of p75<sup>NTR</sup> was determined by SDS-PAGE and immunoblot analysis.  $\beta$ -actin was used as the loading control. C, HuR binds to the p75<sup>NTR</sup> transcript following treatment with R-flurbiprofen or ibuprofen. PC-3 and DU-145 cells were treated for 12 h with 2 mmol/L R-flurbiprofen, 2 mmol/L ibuprofen, or DMSO vehicle control. Cell lysates from equal numbers of cells were incubated with an antibody to HuR, with the exception of the negative controls, and then incubated with protein A/G Plus-agarose beads. Beads were washed, RNA was isolated from the beads, and p75<sup>NTR</sup> mRNA levels were determined by RT-PCR. D, R-flurbiprofen and ibuprofen increase phosphorylation of eIF4E through the p38 MAPK pathway. PC-3 and DU-145 cells were transfected for 72 h with nontargeting siRNA or p38 $\alpha$  siRNA. Following transfection, cells were treated for 30 min with DMSO, 2 mmol/L R-flurbiprofen, or 2 mmol/L ibuprofen. Cell lysates were collected and phosphorylated eIF4E was detected by SDS-PAGE and immunoblot analysis. Blots for phosphorylated eIF4E were stripped and reprobed for total eIF4E.  $\beta$ -actin was used as the loading control.

inhibitor actinomycin D revealed a strong increase in p75<sup>NTR</sup> mRNA stability in cells treated with R-flurbiprofen or ibuprofen. Previously, our investigation of the mechanism by which several prostate cancer cell lines lose expression of p75<sup>NTR</sup> showed that although p75<sup>NTR</sup> is actively transcribed at levels comparable to the high-expressing A875 melanoma cell line, the prostate cancer cells contain <1% of the p75<sup>NTR</sup> mRNA found in A875 cells (10).

Furthermore, transfection of prostate cancer cells with full-length p75<sup>NTR</sup> cDNA led to very modest p75<sup>NTR</sup> protein expression, whereas transfection with p75<sup>NTR</sup> cDNA lacking most of the 3'-UTR resulted in high protein expression. This suggested that although p75<sup>NTR</sup> is transcribed at a high level, prostate cancer cells have very little mRNA or protein due to increased mRNA instability that is mediated through the 3'-UTR. Interestingly, the p75<sup>NTR</sup> 3'-UTR is

~ 2 kb long, which is about four times the length of the average human 3'-UTR (35). It has been suggested that increased 3'-UTR length provides increased potential for posttranscriptional regulation through the 3'-UTR (35). Therefore, it seems likely that R-flurbiprofen and ibuprofen are able to induce p75<sup>NTR</sup> expression by modulating the stability of the transcript and that mRNA stability is an important determinant in regulating the expression level of p75<sup>NTR</sup> protein.

Because induction of p75<sup>NTR</sup> by R-flurbiprofen and ibuprofen treatment seemed to be largely due to increased mRNA stability, we examined activity of the p38 MAPK pathway, which is considered an important regulator of mRNA stability (18, 21, 22). p38 MAPK is activated by phosphorylation, and R-flurbiprofen and ibuprofen caused increased p38 MAPK phosphorylation within 5 min of treatment. Increased activation was still noticeable 8 h following treatment. Because pretreatment with the p38 MAPK selective inhibitor SB202190 or siRNA knockdown of p38 MAPK before profen treatment prevented an induction of p75<sup>NTR</sup>, it seems that the p38 MAPK pathway was involved in regulating profen-mediated induction of p75<sup>NTR</sup>. The data suggest that activation of the p38 MAPK pathway is the primary mechanism responsible for increased p75<sup>NTR</sup> expression. However, inhibition of p38 MAPK did not always result in complete inhibition of p75<sup>NTR</sup> induction, and this may be due to participation of additional pathways. R-Flurbiprofen and ibuprofen induced activation of the kinase MK2, which is directly downstream of p38 MAPK. MK2 and the closely related MK3 are known to be responsible for mediating the mRNA stabilizing effects of the p38 MAPK pathway (22, 23). They are activated with similar kinetics, are able to compensate for one another, have similar substrates, and lead to stabilization of the same group of transcripts in response to lipopolysaccharide (LPS; refs. 23, 25). Not surprisingly, siRNA knockdown of MK2 and MK3 together was more effective in preventing induction of p75<sup>NTR</sup> by R-flurbiprofen or ibuprofen than knockdown of either MK2 or MK3 separately, indicating that p38 MAPK is able to induce p75<sup>NTR</sup> by acting through both MK2 and MK3.

Activation of the p38 MAPK pathway seems to be responsible for R-flurbiprofen- and ibuprofen-mediated induction of p75<sup>NTR</sup> in prostate cancer cells. Significantly, there is an accumulating body of evidence linking p38 MAPK to tumor suppression. For example, inactivation of p38 MAPK, inactivation of the p38 MAPK activating kinases MKK3 and MKK6, and overexpression of the p38 MAPK phosphatase Wip1 are associated with increased tumorigenesis (36, 37). In addition, Wip1 was found to be amplified in primary breast tumors (36, 37). The tumor suppressor p53 is a direct substrate of p38 MAPK and is therefore one way in which p38 MAPK exerts its tumor-suppressing effects (18, 21, 36). However, p53 is inactivated by mutation in roughly half of all cancers (38). This is also the case for PC-3 cells, which are p53-null, and for DU-145 cells, which express mutated p53 (39). Here, we identify induction of p75<sup>NTR</sup> expression as a novel mechanism by which p38 MAPK may achieve tumor suppressor activity. This may be applicable to other cancer types in addition to prostate cancer. For example, we previously showed that ibuprofen up-regulates p75<sup>NTR</sup> in bladder cancer cells, which resulted in p75<sup>NTR</sup>-mediated apoptosis (40). In addition, basic fibroblast growth factor treatment resulted in the p38 MAPK-dependent death of Ewing's sarcoma family of tumor cells and was associated with induction of p75<sup>NTR</sup> that was prevented in the presence of a p38 MAPK selective inhibitor (41). Interestingly, p38 MAPK and p75<sup>NTR</sup> have many overlapping roles in addition to induction of apoptosis and tumor

suppression. For example, both are involved in the regulation of inflammation as well as response to brain injury (18, 24, 42–45). Therefore, it is possible that p38 MAPK achieves these roles, in part, by increasing expression of p75<sup>NTR</sup>. Consistent with this hypothesis is the observation that LPS, which leads to the stabilization of several transcripts through the p38 MAPK/MK2 pathway, also induces p75<sup>NTR</sup> expression (23, 46, 47).

One mechanism by which the p38 MAPK pathway has been shown to stabilize target transcripts involves the RNA binding protein HuR (22, 29–31). HuR is the only RNA binding protein repeatedly shown to stabilize transcripts containing the AUUUA sequence (22, 31). Its ability to stabilize target mRNAs is linked to its subcellular localization, and activation of p38 MAPK and MK2 has been shown to cause translocation of HuR from the nucleus to the cytoplasm, resulting in increased mRNA stability of a number p38 MAPK regulated genes (22, 29–31). The human p75<sup>NTR</sup> transcript contains AUUUA sites located in the 3'-UTR at positions 2,946 and 3,124, and these are conserved in rat and mouse, suggesting that they may be involved in the regulation of p75<sup>NTR</sup> expression. We showed that treatment with R-flurbiprofen or ibuprofen resulted in the binding of HuR to the p75<sup>NTR</sup> transcript, and a modest increase in the cytoplasmic level of HuR was observed following profen treatment. However, siRNA knockdown of HuR before profen treatment only partially prevented an induction of p75<sup>NTR</sup>. These data suggest that binding of HuR to the p75<sup>NTR</sup> transcript is not the sole mechanism responsible for increased p75<sup>NTR</sup> expression. This is not surprising given the elaborate picture of mRNA regulation that is emerging. Recent data suggest that posttranscriptional regulation of gene expression occurs through a plethora of RNA binding proteins that control the splicing, nuclear export, stability, localization, translation, and degradation of transcripts, often through specific sequential or structural elements located in the untranslated regions (48, 49). Therefore, given the exceptionally long 3'-UTR of the p75<sup>NTR</sup> transcript, it is probable that other RNA binding proteins regulate expression as well, and this area requires further investigation. To provide insight into additional mechanisms through which the p38 MAPK pathway may influence p75<sup>NTR</sup> mRNA stability, we examined phosphorylation of eIF4E. eIF4E is phosphorylated by kinases downstream of p38 MAPK (33). It is most well known as a 5'-cap binding protein involved in translation initiation. However, recent data suggest that it regulates gene expression at several levels (34). Interestingly, an increase in eIF4E activity is not associated with an increase in global translation, but rather translation of only a subset of transcripts (34). In addition, eIF4E has been shown to control the nuclear export of a different subset of transcripts (34). Finally, eIF4E is linked to control of mRNA stability in that removal of the 5'-cap is a key step in mRNA degradation, and competition between eIF4E and decapping enzymes has been shown (50). We show that profen treatment increases the level of phosphorylated eIF4E, and this seems to occur, at least partially, through the p38 MAPK pathway because the increase in phosphorylation is substantially inhibited in the presence of p38 MAPK siRNA. Therefore, these data identify modulation of eIF4E activity as another mechanism by which the p38 MAPK pathway may control posttranscriptional events in response to profen treatment.

In this study, we show for the first time that R-flurbiprofen and ibuprofen activate the p38 MAPK pathway in prostate cancer cells and that this pathway is necessary for induction of the p75<sup>NTR</sup> tumor suppressor by these drugs. This mechanism of induction occurs in both androgen-independent PC-3 and DU-145 cells as

well as androgen-responsive LNCaP cells. Retrospective epidemiologic studies have shown that certain nonsteroidal anti-inflammatory drugs, including ibuprofen, reduce the incidence of prostate cancer (15, 16). Numerous recent studies have provided compelling evidence that inhibition of cancer cell growth by nonsteroidal anti-inflammatory drugs can occur through COX-independent mechanisms that have yet to be fully elucidated (12, 13, 17, 40). The enantiomer R-flurbiprofen lacks COX inhibitory activity, indicating that the activation of p38 MAPK and subsequent induction of p75<sup>NTR</sup> is a COX-independent mechanism by which these profens can achieve anticancer activity in the prostate (13). Because p38 MAPK is activated within 5 min of profen treatment, it is likely that R-flurbiprofen and ibuprofen are interacting with signaling

components proximal to p38 MAPK. Further delineation of this mechanism may identify novel therapeutic targets and facilitate the discovery of more effective compounds that are able to induce p75<sup>NTR</sup> expression at lower concentrations.

## Acknowledgments

Received 5/15/2007; revised 9/7/2007; accepted 10/4/2007.

**Grant support:** NIH grant R01DK52626 and the Department of Defense Prostate Cancer Research Program grant PC060409.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We thank Dr. Emil Bogenmann for providing the p75<sup>NTR</sup> promoter-luciferase construct and Dr. Albert J. Fornace for critical reading of the manuscript.

## References

- Chao MV. The p75 neurotrophin receptor. *J Neurobiol* 1994;25:1373–85.
- Chapman BS. A region of the 75 kDa neurotrophin receptor homologous to the death domains of TNFR-I and Fas. *FEBS Lett* 1995;374:216–20.
- Friedman WJ, Greene LA. Neurotrophin signaling via Trks and p75. *Exp Cell Res* 1999;253:131–42.
- Wang X, Bauer JH, Li Y, et al. Characterization of a p75(NTR) apoptotic signaling pathway using a novel cellular model. *J Biol Chem* 2001;276:33812–20.
- Krygiel S, Djakiew D. The neurotrophin receptor p75NTR is a tumor suppressor in human prostate cancer. *Anticancer Res* 2001;21:3749–55.
- Pflug B, Djakiew D. Expression of p75NTR in a human prostate epithelial tumor cell line reduces nerve growth factor-induced cell growth by activation of programmed cell death. *Mol Carcinog* 1998;23:106–14.
- Krygiel S, Djakiew D. Neurotrophin receptor p75(NTR) suppresses growth and nerve growth factor-mediated metastasis of human prostate cancer cells. *Int J Cancer* 2002;98:1–7.
- Khawja F, Tabassum A, Allen J, Djakiew D. The p75(NTR) tumor suppressor induces cell cycle arrest facilitating caspase mediated apoptosis in prostate tumor cells. *Biochem Biophys Res Commun* 2006;341:1184–92.
- Allen J, Khawja F, Djakiew D. Gene therapy of prostate xenograft tumors with a p75NTR lipoplex. *Anticancer Res* 2004;24:2997–3003.
- Krygiel S, Djakiew D. Molecular characterization of the loss of p75(NTR) expression in human prostate tumor cells. *Mol Carcinog* 2001;31:46–55.
- Allen J, Khawja F, Byers S, Djakiew D. The p75NTR mediates a bifurcated signal transduction cascade through the NFκB and JNK pathways to inhibit cell survival. *Exp Cell Res* 2005;304:69–80.
- Tegeder I, Pfeilschifter J, Geisslinger G. Cyclooxygenase-independent actions of cyclooxygenase inhibitors. *FASEB J* 2001;15:2057–72.
- Wechter WJ, Leipold DD, Murray ED, Jr., et al. E-7869 (R-flurbiprofen) inhibits progression of prostate cancer in the TRAMP mouse. *Cancer Res* 2000;60:2203–8.
- Andrews J, Djakiew D, Krygiel S, Andrews P. Superior effectiveness of ibuprofen compared with other NSAIDs for reducing the survival of human prostate cancer cells. *Cancer Chemother Pharmacol* 2002;50:277–84.
- Nelson JE, Harris RE. Inverse association of prostate cancer and non-steroidal anti-inflammatory drugs (NSAIDs): results of a case-control study. *Oncol Rep* 2000;7:169–70.
- Harris RE, Beebe-Donk J, Doss H, Burr Doss D. Aspirin, ibuprofen, and other non-steroidal anti-inflammatory drugs in cancer prevention: a critical review of non-selective COX-2 blockade (review). *Oncol Rep* 2005;13:559–83.
- Quann EJ, Khawja F, Zavitz KH, Djakiew D. The aryl propionic acid R-flurbiprofen selectively induces p75NTR-dependent decreased survival of prostate tumor cells. *Cancer Res* 2007;67:3254–62.
- Zarubín T, Han J. Activation and signaling of the p38 MAP kinase pathway. *Cell Res* 2005;15:11–8.
- Olson JM, Hallahan AR. p38 MAP kinase: a convergence point in cancer therapy. *Trends Mol Med* 2004;10:125–9.
- Kennedy NJ, Cellurale C, Davis RJ. A radical role for p38 MAPK in tumor initiation. *Cancer Cell* 2007;11:101–3.
- Gaestel M. MAPKAP kinases—MKs—two's company, three's a crowd. *Nat Rev Mol Cell Biol* 2006;7:120–30.
- Tran H, Maurer F, Nagamine Y. Stabilization of urokinase and urokinase receptor mRNAs by HuR is linked to its cytoplasmic accumulation induced by activated mitogen-activated protein kinase-activated protein kinase 2. *Mol Cell Biol* 2003;23:7177–88.
- Ronkina N, Kotlyarov A, Ttrich-Breiholz O, et al. The mitogen-activated protein kinase (MAPK)-activated protein kinases MK2 and MK3 cooperate in stimulation of tumor necrosis factor biosynthesis and stabilization of p38 MAPK. *Mol Cell Biol* 2007;27:170–81.
- Peterson S, Bogenmann E. Osmotic swelling induces p75 neurotrophin receptor (p75NTR) expression via nitric oxide. *J Biol Chem* 2003;278:33943–50.
- Clifton AD, Young PR, Cohen P. A comparison of the substrate specificity of MAPKAP kinase-2 and MAPKAP kinase-3 and their activation by cytokines and cellular stress. *FEBS Lett* 1996;392:209–14.
- Brennan CM, Steitz JA. HuR and mRNA stability. *Cell Mol Life Sci* 2001;58:266–77.
- Chen CY, Xu N, Shyu AB. Highly selective actions of HuR in antagonizing AU-rich element-mediated mRNA destabilization. *Mol Cell Biol* 2002;22:7268–78.
- Subbaramaiah K, Marmo TP, Dixon DA, Dannenberg AJ. Regulation of cyclooxygenase-2 mRNA stability by taxanes: evidence for involvement of p38, MAPKAPK-2, and HuR. *J Biol Chem* 2003;278:37637–47.
- Lin FY, Chen YH, Lin YW, et al. The role of human antigen R, an RNA-binding protein, in mediating the stabilization of toll-like receptor 4 mRNA induced by endotoxin: a novel mechanism involved in vascular inflammation. *Arterioscler Thromb Vasc Biol* 2006;26:2622–9.
- Jin SH, Kim TI, Yang KM, Kim WH. Thalidomide destabilizes cyclooxygenase-2 mRNA by inhibiting p38 mitogen-activated protein kinase and cytoplasmic shuttling of HuR. *Eur J Pharmacol* 2007;558:14–20.
- Song IS, Tatebe S, Dai W, Kuo MT. Delayed mechanism for induction of γ-glutamylcysteine synthetase heavy subunit mRNA stability by oxidative stress involving p38 mitogen-activated protein kinase signaling. *J Biol Chem* 2005;280:28230–40.
- Pascale A, Amadio M, Scapagnini G, et al. Neuronal ELAV proteins enhance mRNA stability by a PKCα-dependent pathway. *Proc Natl Acad Sci U S A* 2005;102:12065–70.
- Scheper GC, Proud CG. Does phosphorylation of the cap-binding protein eIF4E play a role in translation initiation? *Eur J Biochem* 2002;269:5350–9.
- Culjkovic B, Topisirovic I, Borden KL. Controlling gene expression through RNA regulons: the role of the eukaryotic translation initiation factor eIF4E. *Cell Cycle* 2007;6:65–9.
- Mazumder B, Seshadri V, Fox PL. Translational control by the 3'-UTR: the ends specify the means. *Trends Biochem Sci* 2003;28:91–8.
- Bulavin DV, Demidov ON, Saito S, et al. Amplification of PPM1D in human tumors abrogates p53 tumor-suppressor activity. *Nat Genet* 2002;31:210–5.
- Brancho D, Tanaka N, Jaeschke A, et al. Mechanism of p38 MAP kinase activation *in vivo*. *Genes Dev* 2003;17:1969–78.
- Joerger AC, Fersht AR. Structure-function-rescue: the diverse nature of common p53 cancer mutants. *Oncogene* 2007;26:2226–42.
- Mori A, Lehmann S, O'Kelly J, et al. Capsaicin, a component of red peppers, inhibits the growth of androgen-independent, p53 mutant prostate cancer cells. *Cancer Res* 2006;66:3222–9.
- Khawja F, Allen J, Lynch J, Andrews P, Djakiew D. Ibuprofen inhibits survival of bladder cancer cells by induced expression of the p75NTR tumor suppressor protein. *Cancer Res* 2004;64:6207–13.
- Williamson AJ, Dibling BC, Boyne JR, Selby P, Burchill SA. Basic fibroblast growth factor-induced cell death is effected through sustained activation of p38MAPK and up-regulation of the death receptor p75NTR. *J Biol Chem* 2004;279:47912–28.
- Wang X, Xu L, Wang H, Young PR, Gaestel M, Feuerstein GZ. Mitogen-activated protein kinase-activated protein (MAPKAP) kinase 2 deficiency protects brain from ischemic injury in mice. *J Biol Chem* 2002;277:43968–72.
- Nassenstein C, Kerzel S, Braun A. Neurotrophins and neurotrophin receptors in allergic asthma. *Prog Brain Res* 2004;146:347–67.
- Raychaudhuri SP, Raychaudhuri SK. Role of NGF and neurogenic inflammation in the pathogenesis of psoriasis. *Prog Brain Res* 2004;146:433–7.
- Chien CC, Fu WM, Huang HI, et al. Expression of neurotrophic factors in neonatal rats after peripheral inflammation. *J Pain* 2007;8:161–7.
- Hennigan A, Trotter C, Kelly AM. Lipopolysaccharide impairs long-term potentiation and recognition memory and increases p75NTR expression in the rat dentate gyrus. *Brain Res* 2007;1130:158–66.
- Watt JA, Paden CM. Up-regulation of the p75 low-affinity neurotrophin receptor by phagocytically active perivascular active cells in the rat neural lobe. *Cell Tissue Res* 2001;303:81–91.
- Moore MJ. From birth to death: the complex lives of eukaryotic mRNAs. *Science* 2005;309:1514–8.
- Keene JD. RNA regulons: coordination of post-transcriptional events. *Nat Rev Genet* 2007;8:533–43.
- von der Haar T, Gross JD, Wagner G, McCarthy JE. The mRNA cap-binding protein eIF4E in post-transcriptional gene expression. *Nat Struct Mol Biol* 2004;11:503–11.

# The Aryl Propionic Acid *R*-Flurbiprofen Selectively Induces p75<sup>NTR</sup>-Dependent Decreased Survival of Prostate Tumor Cells

Emily J. Quann,<sup>1</sup> Fatima Khwaja,<sup>1</sup> Kenton H. Zavitz,<sup>3</sup> and Daniel Djakiew<sup>1,2</sup>

<sup>1</sup>Department of Biochemistry and Molecular Cellular Biology and the <sup>2</sup>Vincent T. Lombardi Cancer Center, Georgetown University Medical Center, Washington, DC, and <sup>3</sup>Myriad Pharmaceuticals Inc., Salt Lake City, Utah

## Abstract

Epidemiologic studies show that patients chronically consuming nonsteroidal anti-inflammatory drugs (NSAID) for arthritis exhibit a reduced incidence of prostate cancer. In addition, some NSAIDs show anticancer activity *in vitro*. NSAIDs exert their anti-inflammatory effects by inhibiting cyclooxygenase (COX) activity; however, evidence suggests that COX-independent mechanisms mediate decreased prostate cancer cell survival. Hence, we examined the effect of selected aryl propionic acid NSAIDs and structurally related compounds on the decreased survival of prostate cancer cell lines PC-3, DU-145, and LNCaP by induction of the p75<sup>NTR</sup> protein. The p75<sup>NTR</sup> has been shown to function as a tumor suppressor in the prostate by virtue of its intracellular death domain that can initiate apoptosis and inhibit growth. The most efficacious compounds for induction of p75<sup>NTR</sup> and decreased survival, in rank-order, were *R*-flurbiprofen, ibuprofen, oxaprozin, fenoprofen, naproxen, and ketoprofen. Because *R*-flurbiprofen and ibuprofen exhibited the greatest efficacy, we examined their dose-dependent specificity of induction for p75<sup>NTR</sup> relative to other members of the death receptor family. Whereas treatment with *R*-flurbiprofen or ibuprofen resulted in a massive induction of p75<sup>NTR</sup> protein levels, the expression of Fas, p55<sup>TNFR</sup>, DR3, DR4, DR5, and DR6 remained largely unchanged. Moreover, transfection of either cell line before *R*-flurbiprofen or ibuprofen treatment with a dominant negative form of p75<sup>NTR</sup> to antagonize p75<sup>NTR</sup> activity or p75<sup>NTR</sup> small interfering RNA to prevent p75<sup>NTR</sup> protein expression rescued both cell lines from decreased survival. Hence, *R*-flurbiprofen and ibuprofen selectively induce p75<sup>NTR</sup>-dependent decreased survival of prostate cancer cells independently of COX inhibition. [Cancer Res 2007;67(7):1–9]

## Introduction

Prostate cancer is the most commonly diagnosed cancer and ranks as the second leading cause of cancer-related deaths among men in the United States (1, 2). Recent epidemiologic studies found a correlation between long-term nonsteroidal anti-inflammatory drug (NSAID) use and decreased prostate cancer risk (3–6). In addition, many *in vitro* studies involving various human prostate cancer cell lines consistently show decreased proliferation and increased apoptosis in response to select NSAID treatment (7, 8). NSAIDs exert their anti-inflammatory activity by inhibiting cyclo-

oxygenase (COX), the enzyme which catalyzes the conversion of arachidonic acid to prostaglandins. Two isoforms of COX exist; COX-1 is a housekeeping gene that is constitutively expressed at low levels in most cells types, whereas COX-2 is highly inducible in response to cytokines, hormones, and growth factors. COX-2 seems to play a significant role in the promotion of colon cancer with 50% of precancerous adenomatous polyps and 85% of colon carcinomas exhibiting COX-2 overexpression (9). However, the data pertaining to the role of COX-2 in prostate cancer are less conclusive. Although some studies show overexpression, others show expression is low or absent relative to normal tissue (10–14). In addition, there is not a consensus regarding the status of COX-2 expression in established prostate cancer cell lines, including LNCaP, DU-145, and PC-3 (13, 15–17). However, regardless of COX-2 expression, these cell lines all show susceptibility to select NSAID treatment (17–19). Furthermore, the anticancer activity of NSAIDs occurs at concentrations several orders of magnitude higher than those required to inhibit COX-2, and different NSAIDs show varying levels of anticancer activity (20–22). These results suggest the existence of a COX-independent mechanism by which some NSAIDs initiate apoptosis and inhibit proliferation.


The p75<sup>NTR</sup> (neurotrophin receptor) is a member of the tumor necrosis factor receptor (TNFR) superfamily capable of inducing apoptosis through a conserved intracellular death domain (23, 24). It binds neurotrophin ligands with similar affinity; however, unlike other members of the TNFR superfamily, p75<sup>NTR</sup> induces cell death and suppresses growth in the unbound state (25–29). Recently, p75<sup>NTR</sup> was identified as a tumor and metastasis suppressor in the prostate and bladder (27, 28). Although normal prostate epithelial cells express high levels of p75<sup>NTR</sup>, this expression becomes suppressed as prostate cancer progresses (30). In addition, the human prostate cancer cell lines PC-3, DU-145, and LNCaP, all derived from metastases, show little to no p75<sup>NTR</sup> expression (31). However, the p75<sup>NTR</sup> gene in these cells has remained intact, indicating that the potential for up-regulation may exist (31). Furthermore, exogenous reexpression of p75<sup>NTR</sup> in PC-3 cells suppressed growth and increased apoptosis (28, 32). This suggests potential for treatments that result in the up-regulation and reexpression of p75<sup>NTR</sup> in prostate cancer cells.

Ibuprofen and flurbiprofen belong to the aryl propionic acid class of NSAIDs. Treatment of T24 bladder cancer cells and HCT-116 colon cancer cells with ibuprofen or the enantiomer *R*-flurbiprofen, which lacks COX inhibitory activity, induced a dose-dependent up-regulation of p75<sup>NTR</sup> and a corresponding decrease in survival (33). Rescue experiments involving transfection of dominant negative forms of p75<sup>NTR</sup> before ibuprofen treatment of T24 cells showed that the observed induction of p75<sup>NTR</sup> was causal of the decreased survival (33). Significantly, *R*-flurbiprofen treatment of TRAMP mice has been shown to lower the incidence of primary tumors and metastases of prostate cancer (34). In this context, we show that *R*-flurbiprofen and ibuprofen most

**Requests for reprints:** Daniel Djakiew, Department of Biochemistry and Molecular & Cellular Biology, Georgetown University Medical Center, 3900 Reservoir Road, N.W., Washington, DC 20057-1436. Phone: 202-687-1203; Fax: 1-202-687-1823; E-mail: djakiewd@georgetown.edu.

©2007 American Association for Cancer Research.  
doi:10.1158/0008-5472.CAN-06-3657





**Figure 1.** A, expression of the p75<sup>NTR</sup> in PC-3 and DU-145 cells following 48-h treatment with 0, 0.1, 0.25, 0.5, 1.0, or 2.0 mmol/L *R*-flurbiprofen, ibuprofen, oxaprozin, fenoprofen, naproxen, or ketoprofen. Cell lysates were collected, and 50 µg of protein were subjected to SDS-PAGE and immunoblot analysis using a monoclonal antibody to human p75<sup>NTR</sup>. The A875 melanoma cell line was used as a positive control (+ve). B, expression of p75<sup>NTR</sup> in LNCaP cells following 48-h treatment with 0 to 2 mmol/L *R*-flurbiprofen or ibuprofen. SDS-PAGE was done using 50 µg of protein followed by immunoblot analysis. C, p75<sup>NTR</sup> and GAPDH mRNA levels in PC-3 and DU-145 cells following 0, 2, 4, 8, 12, 18, and 24 h of 2.0 mmol/L *R*-flurbiprofen or ibuprofen treatment determined by RT-PCR. D, expression of COX-1 and COX-2 in PC-3 and DU-145 cells following 48-h treatment with 0, 0.1, 0.25, 0.5, 1.0, or 2.0 mmol/L *R*-flurbiprofen or ibuprofen. SDS-PAGE was done using cell lysates containing 50 µg of protein and followed by immunoblot analysis using a monoclonal antibody to human COX-1 and goat polyclonal antibody to human COX-2. U-937 and RAW 264.7 cell lysates were used as positive controls for COX-1 and COX-2, respectively.

effectively up-regulate p75<sup>NTR</sup> expression in both PC-3 and DU-145 cell lines relative to other aryl propionic acids. This up-regulation occurs in a dose-dependent manner and corresponds to a decrease in cell survival. The use of p75<sup>NTR</sup> dominant negatives and p75<sup>NTR</sup>-targeting small interfering RNA (siRNA) shows that the observed decreased survival is directly mediated through p75<sup>NTR</sup>, a COX-independent mechanism.

## Materials and Methods

**Cell lines and culture conditions.** PC-3 and DU-145 cell lines were obtained from the tissue culture core facility of the Georgetown University Lombardi Comprehensive Cancer Center. LNCaP cells were obtained from the American Type Culture Collection (Rockville, MD). All cell lines were

maintained in DMEM (Mediatech Inc., Herndon, VA) containing 4.5 g/L glucose and L-glutamine supplemented with antibiotic/antimycotic [100 units/mL penicillin G, 100 µg/mL streptomycin, and 0.25 µg/mL amphotericin B (Mediatech Inc.)] and 5% fetal bovine calf serum (Sigma Chemical Co., St. Louis, MO). Cells were incubated in the presence of 5% CO<sub>2</sub> and air at 37°C.

**Drug preparation, treatment, and cell lysis.** Stock solutions were prepared by dissolving each aryl propionic acid [ibuprofen, ketoprofen, naproxen, oxaprozin, fenoprofen (Sigma), and *R*-flurbiprofen (Myriad Pharmaceuticals Inc., Salt Lake City, UT)] in DMSO (Sigma) at a concentration of 200 mmol/L. Cells were seeded overnight at 70% to 80% confluency and were then treated for 48 h at concentrations of 0, 0.1, 0.25, 0.5, 1.0, and 2.0 mmol/L. Cell lysates of treated cells were prepared using lysis buffer [10 mmol/L Tris-Cl (pH, 7.4), 10 mmol/L NaCl, 3 mmol/L MgCl<sub>2</sub>, and 0.5% Nonidet P-40] containing 1 µL/mL cocktail protease inhibitor


(Sigma). The supernatant was retained, and protein concentration was determined according to the manufacturer's protocol (Bio-Rad Laboratories, Hercules, CA). For long-term treatment of cells, ~1000 cells were seeded overnight and treated the next day with 0, 0.1, 0.2, 0.3, 0.4, and 0.5 mmol/L ibuprofen or *R*-flurbiprofen twice per week for 2 weeks. After 2 weeks, lysates were prepared as described above, or cells were trypsinized, resuspended, and counted using a hemocytometer.

**Immunoblot analysis.** Immunoblot analysis was done by loading 50 µg of protein onto 10% SDS-polyacrylamide gels for electrophoresis, followed by transfer to a nitrocellulose membrane (Amersham Pharmacia Biotech, Piscataway, NJ). Membranes were blocked in 5% nonfat milk (Bio-Rad Laboratories) and then incubated in the primary antibody: murine monoclonal anti-p75<sup>NTR</sup> (1:2,000, Upstate Cell Signaling Solutions, Lake Placid, NY), murine monoclonal anti-Fas (1:100, Santa Cruz Biotechnologies, Santa Cruz, CA), murine monoclonal anti-p55<sup>TNFR</sup> (1:100, Santa Cruz Biotechnologies), rat polyclonal anti-DR3 (1:200, Santa Cruz Biotechnologies), murine monoclonal anti-DR4 (1:100, Santa Cruz Biotechnologies), rat polyclonal anti-DR5 (1:500, ProSci Inc., Poway, CA), rat polyclonal anti-DR6 (2 µg/mL; Upstate Cell Signaling Solutions), mouse monoclonal anti-COX-1 (1:200, Santa Cruz Biotechnologies), goat polyclonal anti-COX-2 (1:200, Santa Cruz Biotechnologies), or murine monoclonal anti-β-actin (1:5000, Sigma). Following incubation in the primary antibody, membranes were incubated in the appropriate horseradish peroxidase-conjugated secondary antibody (1:3,000, Bio-Rad Laboratories). Immunoreactivity was detected using the chemiluminescence detection reagent (Amersham Pharmacia Biotech). Several positive controls were used: A875 whole cell lysate (WCL; Tissue Culture Core Facility, Georgetown University) for p75<sup>NTR</sup>, A431 WCL (Santa Cruz Biotechnologies) for Fas, MCF-7 WCL (Santa Cruz Biotechnologies) for p55<sup>TNFR</sup>, Jurkat WCL (Upstate Cell Signaling Solutions) for DR3

and DR6, HeLa WCL (ProSci Inc.) for DR4 and DR5, U-937 WCL (Santa Cruz Biotechnologies) for COX-1, and RAW 264.7 WCL (Santa Cruz Biotechnologies) for COX-2.

**Reverse transcription-PCR.** PC-3 and DU145 cells were treated with 2.0 mmol/L *R*-flurbiprofen or ibuprofen. RNA was isolated following 0, 2, 4, 8, 12, 18, and 24 h of treatment using TRIzol Reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol. Reverse transcription-PCR (RT-PCR) was done using the SuperScript III One-Step RT-PCR System with Platinum Taq DNA Polymerase (Invitrogen) using 250 ng RNA for p75<sup>NTR</sup> and 125 ng RNA for glyceraldehyde-3-phosphate dehydrogenase (GAPDH). Primers were designed using Primer Quest, and their sequences are as follows: p75<sup>NTR</sup> forward 5'-AGG TGA CCT TCT GGG AAA TGG CTT-3', p75<sup>NTR</sup> reverse 5'-ATT TCC TCC GAT GCT TCT CTG GCA-3', GAPDH forward 5'-CCA CCC ATG GCA AAT TCC ATG GCA-3', and GAPDH reverse 5'-TCT AGA CGG CAG GTC AGG TCC ACC-3' (Integrated DNA Technologies, Coralville, IA). cDNA synthesis was done at 47°C for 30 min followed by denaturation at 94°C for 2 min and then 30 cycles of PCR at 94°C for 1 min, 60°C for 1 min, and 72°C for 1 min, with final extension at 72°C for 5 min. PCR products were separated on 1.5% agarose gels.

**Hoechst dye nuclear (DNA) staining.** Hoechst staining to identify apoptotic nuclei was conducted as described previously (33). PC-3, DU-145, and LNCaP cells were treated for 48 h with aryl propionic acids and then fixed in 10% formalin (Electron Microscopy Sciences, Hatfield, PA). Cells were collected, washed in PBS, centrifuged, resuspended in PBS, and dried on slides. Slides were rehydrated with PBS, washed with distilled water, and covered with a 1:1,000 aqueous dilution of Hoechst 33258 stain (Molecular Probes, Eugene, OR) for a final concentration of 10 µg/mL. After 10 min, slides were washed with distilled water, mounted, and viewed using a fluorescence microscope (Zeiss Axioplan 2 Imaging, Jena, Germany).



**Figure 2.** A and B, PC-3, DU-145, and LNCaP cell survival analysis by MTT assay following treatment with 0, 0.1, 0.25, 0.5, 1.0, or 2.0 mmol/L *R*-flurbiprofen or ibuprofen for 48 h. Columns, mean relative to the control (0 mM); bars, SE. \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ . C and D, detection of apoptotic nuclei (arrows) by Hoechst staining of PC-3, DU-145, and LNCaP cells following treatment with 0 or 2.0 mmol/L *R*-flurbiprofen or ibuprofen.



**p75<sup>NTR</sup> dominant negative transfection.** PC-3 and DU-145 cells were transiently transfected with one of two p75<sup>NTR</sup> dominant negative vectors described previously (35). The  $\Delta$ DD vector expresses p75<sup>NTR</sup> with the death domain deleted, and the  $\Delta$ ICD vector expresses p75<sup>NTR</sup> with the slightly larger intracellular domain deleted. Both are ecdysone-inducible p75<sup>NTR</sup> vectors and, therefore, were each cotransfected with the ecdysone receptor plasmid pVgRxR. The transfection was done with LipofectAMINE reagent (Invitrogen) in serum-free medium for 6 h, after which serum-containing medium was added. After 18 subsequent hours, cells were incubated in 1  $\mu$ mol/L ponasterone A (Invitrogen) for 24 h to drive the expression of the dominant negative gene products. Following incubation with ponasterone A, cells were treated with aryl propionic acids for 48 h, and relative cell survival was determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay (Roche Applied Science, Indianapolis, IN).


**siRNA transfection.** PC-3 and DU-145 cells were transfected for 24 h with nontargeting or p75<sup>NTR</sup>-specific siRNA [Dharmacon RNA Technologies (Duplex D-009340-03), Lafayette, CO] at a final concentration of 100 nmol/L according to the manufacturer's protocol using transfection reagents DharmaFECT 1 for DU-145 cells and DharmaFECT 2 (Dharmacon RNA Technologies) for PC-3 cells. After transfection, the cells were treated with aryl propionic acids for 48 h, followed by determination of p75<sup>NTR</sup> protein expression or relative cell survival by MTT assay (Roche Applied Science).

## Results

**Aryl propionic acids selectively induce p75<sup>NTR</sup> expression and decrease cell survival.** We examined the ability of *R*-flurbiprofen and ibuprofen, as well as four other members of the aryl propionic acid family, including oxaprozin, fenoprofen, naproxen, and ketoprofen to induce the expression of p75<sup>NTR</sup> protein in PC-3 and DU-145 human prostate cancer cells. In both cell lines, *R*-flurbiprofen was the most efficacious for the induction of p75<sup>NTR</sup> expression followed by ibuprofen, oxaprozin, fenoprofen, naproxen, and finally, ketoprofen, which was the least effective compound (Fig. 1A). For subsequent experiments, only *R*-flurbiprofen and ibuprofen were used because they were the most effective of the six aryl propionic acids tested. Treatment with *R*-flurbiprofen or ibuprofen also resulted in an induction of p75<sup>NTR</sup> in LNCaP cells, which are androgen responsive (Fig. 1B). To elucidate the mechanism by which these aryl propionic acids may be inducing p75<sup>NTR</sup> protein expression, we determined relative p75<sup>NTR</sup> mRNA levels at various time points following 2.0 mmol/L *R*-flurbiprofen or ibuprofen treatment (Fig. 1C). In both cell lines, *R*-flurbiprofen and ibuprofen significantly induced p75<sup>NTR</sup> message level. This induction is first noticeable within 4 h of treatment in both cell lines with both drugs.

The aryl propionic acids are commonly used as NSAIDs that act through COX inhibition. Therefore, we examined the status of COX expression in both cell lines and determined whether the expression level of either isoform changed in response to treatment with *R*-flurbiprofen and ibuprofen (Fig. 1D). Both PC-3 and DU-145 cells expressed COX-1 at low levels, and expression remained unchanged following *R*-flurbiprofen or ibuprofen treatment. PC-3 cells lacked COX-2 expression; however, expression was induced upon treatment with *R*-flurbiprofen or ibuprofen. DU-145 cells also lacked COX-2 expression, and no induction was observed following treatment with either compound.

Examination of the effect of *R*-flurbiprofen and ibuprofen on survival of PC-3, DU-145, and LNCaP cells after 48-h treatment resulted in a dose-dependent decrease in survival that corresponded with the observed induction of p75<sup>NTR</sup> (Fig. 2A and B). Again, *R*-flurbiprofen was more efficacious than ibuprofen, resulting in a greater dose-dependent decrease in survival. Because



**Figure 3.** A and B, expression of TNFR superfamily members in PC-3 cells after 48-h treatment with 0, 0.1, 0.25, 0.5, 1.0, or 2.0 mmol/L *R*-flurbiprofen or ibuprofen. About 50  $\mu$ g of cell lysate were subjected to SDS-PAGE followed by immunoblot analysis using monoclonal antibodies to human Fas, p55<sup>TNFR</sup>, and DR3, or rat polyclonal antibodies to human DR3, DR5, and DR6, and  $\beta$ -actin as the loading control. A431, MCF-7, Jurkat, and HeLa cell lysates were used as positive controls for Fas, p55<sup>TNFR</sup>, DR3 and DR6, and DR4 and DR5, respectively.

p75<sup>NTR</sup> contains an intracellular death domain capable of initiating apoptosis, we used Hoechst staining to identify fragmented nuclei typical of apoptotic cells. Treatment with 2.0 mmol/L *R*-flurbiprofen or ibuprofen resulted in an induction of apoptotic cells, indicating that apoptosis was at least partially responsible for the observed decrease in survival (Fig. 2C and D).


p75<sup>NTR</sup> is a member of the TNFR superfamily in which members contain an intracellular death domain capable of inducing apoptosis (23, 24). Although 48-h treatment with *R*-flurbiprofen resulted in a substantial dose-dependent induction of p75<sup>NTR</sup>, the other death receptors, including Fas, p55<sup>TNFR</sup>, DR3, DR4, DR5, and DR6, showed little, if any, induction in PC-3 cells (Fig. 3A). Similar results were observed following treatment with ibuprofen (Fig. 3B). There seemed to be a slight induction of DR4 with ibuprofen; however, the magnitude of the response was small relative to the induction observed for p75<sup>NTR</sup> expression (Fig. 3B). The addition of the ligand corresponding to each death receptor had no effect on the induction of death receptors by *R*-flurbiprofen or ibuprofen (data not shown). Therefore, *R*-flurbiprofen and ibuprofen selectively induced p75<sup>NTR</sup>, whereas protein expression of the other TNFR family members remained largely unchanged.

**Long-term R-flurbiprofen and ibuprofen treatment results in p75<sup>NTR</sup> induction and decreased survival at lower drug concentrations.** Epidemiologic studies have shown a correlation between long-term continuous NSAID use and a decreased risk of prostate cancer (3, 4). Therefore, we examined whether chronic treatment with R-flurbiprofen or ibuprofen for 2 weeks at lower concentrations of 0, 0.1, 0.2, 0.3, 0.4, and 0.5 mmol/L may be equally effective or more effective than the 48-h acute treatment at higher concentrations in PC-3 and DU-145 cells. Long-term treatment with R-flurbiprofen resulted in a dose-dependent decrease in cell growth with severe growth arrest even at 0.1 mmol/L and almost a complete loss of growth at 0.3 mmol/L for PC-3 cells and 0.5 mmol/L for DU-145 cells (Fig. 4A). In addition, there was a dose-dependent increase in p75<sup>NTR</sup> levels beginning at 0.1 mmol/L, which corresponds to the concentration at which decreased growth first occurred (Fig. 4B). Similarly, long-term treatment with ibuprofen resulted in a dose-dependent decrease in survival and induction of p75<sup>NTR</sup> at lower concentrations than observed with 48-h ibuprofen treatment, but was not as efficacious as long-term treatment with R-flurbiprofen (Fig. 4C and D).


**Decreased survival caused by R-flurbiprofen and ibuprofen is mediated through p75<sup>NTR</sup>.** The results presented thus far show that the induction of p75<sup>NTR</sup> is associated with decreased survival of PC-3 and DU-145 cells in response to R-flurbiprofen or ibuprofen treatment. To determine if the observed decreased survival is causally mediated through R-flurbiprofen- and ibuprofen-dependent induction of p75<sup>NTR</sup>, ponasterone A-inducible expression vectors for one of two different dominant negative forms of p75<sup>NTR</sup> were transfected into PC-3 and DU-145 cells before R-flurbiprofen or ibuprofen treatment. The dominant negative vectors both express truncated forms of p75<sup>NTR</sup>.  $\Delta$ DDp75<sup>NTR</sup> has a deletion of the death

domain, and  $\Delta$ ICDp75<sup>NTR</sup> has a deletion of the larger intracellular domain that includes the death domain. PC-3 and DU-145 cells treated with 0, 0.5, or 1.0 mmol/L R-flurbiprofen alone or with ponasterone A resulted in a similar dose-dependent decrease in survival (Fig. 5A and B). R-flurbiprofen treatment of cells transfected with either  $\Delta$ DDp75<sup>NTR</sup> or  $\Delta$ ICDp75<sup>NTR</sup> but not treated with ponasterone A yielded the same result (Fig. 5A and B). However, cells transfected with either of the two dominant negative expression vectors followed by ponasterone A treatment to induce expression of the gene product exhibited increased survival following R-flurbiprofen treatment compared with cells not expressing dominant negative p75<sup>NTR</sup> (Fig. 5A and B). Similar results were observed with ibuprofen in PC-3 and DU-145 cells, where expression of the dominant negative forms of p75<sup>NTR</sup> partially prevented decreased survival due to ibuprofen (Fig. 5C and D).

We did similar rescue experiments using siRNA targeted against p75<sup>NTR</sup>. Transfection of PC-3 cells with p75<sup>NTR</sup> siRNA before R-flurbiprofen almost completely prevented the induction of p75<sup>NTR</sup> protein, whereas transfection with nontargeting siRNA did not prevent the induction of p75<sup>NTR</sup> protein (Fig. 6A). Treatment with R-flurbiprofen at 0, 0.5, and 1.0 mmol/L in untransfected PC-3 cells or in PC-3 cells transfected with nontargeting siRNA resulted in a similar dose-dependent decrease in cell viability (Fig. 6A). However, treatment of PC-3 cells transfected with p75<sup>NTR</sup> siRNA partially prevented R-flurbiprofen-dependent decreased survival at 0.5 and 1.0 mmol/L concentrations (Fig. 6A). Similarly, transfection of PC-3 or DU-145 cells with p75<sup>NTR</sup>-targeting siRNA before ibuprofen treatment also almost completely blocked the induction of p75<sup>NTR</sup>, whereas transfection with nontargeting siRNA did not block induction (Fig. 6C and D). Treatment with 0, 0.5, and 1.0 mmol/L ibuprofen led to a dose-dependent decrease of cell survival in



**Figure 4.** A and C, relative growth of PC-3 and DU-145 cells following treatment with 0, 0.1, 0.2, 0.3, 0.4, or 0.5 mmol/L R-flurbiprofen or ibuprofen twice per week for 2 wks. Cell number was determined using a hemocytometer. Columns, mean relative to the control (0 mM); bars, SE. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ . B and D, expression of p75<sup>NTR</sup> in PC-3 and DU-145 cells that were treated with 0, 0.1, 0.2, 0.3, 0.4, or 0.5 mmol/L R-flurbiprofen or ibuprofen twice per week for 2 wks. The A875 cell line was used as a positive control. SDS-PAGE was done using 50  $\mu$ g of cell lysate and followed by immunoblot analysis using a monoclonal antibody to human p75<sup>NTR</sup> and  $\beta$ -actin as the loading control.



**Figure 5.** A to D, PC-3 and DU-145 cell survival analysis by MTT assay following 48-h treatment with 0, 0.5, or 1.0 mmol/L *R*-flurbiprofen or ibuprofen. Before treatment, cells were cotransfected for 6 h in serum-free medium with ponasterone A-inducible ecdysone receptor plasmid pVgRxR and either  $\Delta$ DDp75<sup>NTR</sup> or  $\Delta$ ICDp75<sup>NTR</sup>, which express a truncated dominant negative p75<sup>NTR</sup> protein that lacks either the death domain or the entire intracellular domain, respectively. Following transfection, cells were incubated in serum-containing medium for 18 h and then incubated in 1  $\mu$ mol/L ponasterone A (P) for 24 h to drive expression of the dominant negative gene products. Columns, mean relative to the control (0 mM); bars, SE. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ .

untransfected cells and in cells transfected with nontargeting siRNA, whereas transfection with p75<sup>NTR</sup> siRNA before ibuprofen treatment prevented a decrease in survival at the 1.0 mmol/L concentration (Fig. 6C and D). Transfection of DU-145 cells with p75<sup>NTR</sup>-targeting siRNA only partially prevented p75<sup>NTR</sup> induction upon *R*-flurbiprofen treatment (Fig. 6B). Consistently, only a partial rescue from *R*-flurbiprofen-dependent decreased survival was observed in these cells.

## Discussion

The profens, also referred to as 2-aryl propionic acid derivatives, are a class of NSAIDs that share several characteristics. They can be reversible inhibitors of COX, are highly bound to plasma albumin, and are weak acids (36). They exist as enantiomer pairs, and it is generally the *S*-enantiomer, but not the *R*-enantiomer, that possesses potent COX inhibitory activity (37). Some profens have been in use for roughly 30 years, most commonly as treatment for inflammation due to rheumatoid arthritis (38–43). However, based on emerging studies associating NSAIDs with anticancer activity, various profens have been examined for their efficacy as chemopreventive and chemotherapeutic agents in a variety of cancer types (44–46). In this study, we examined the ability of six profens, including *R*-flurbiprofen, ibuprofen, oxaprozin, fenoprofen, naproxen, and ketoprofen, to induce the expression of the p75<sup>NTR</sup>

tumor suppressor in prostate cancer cells. We observed a dose-dependent induction of p75<sup>NTR</sup> by these drugs; however, the level of efficacy varied with *R*-flurbiprofen followed by ibuprofen as the most effective. Consistently, previous reports identify these two drugs as promising agents in prostate cancer treatment. *R*-flurbiprofen was shown to inhibit the progression of prostate cancer in the TRAMP mouse, whereas ibuprofen was shown to reduce survival of LNCaP and DU-145 human prostate cancer cells (8, 21, 34). In addition, treatment of the human colon cancer COX-null cell line HCT-116 with either of these drugs resulted in reduced cell survival, indicating that these drugs possess the ability to inhibit growth through a COX-independent mechanism (33). The remaining four profens induced p75<sup>NTR</sup> to a lesser degree with oxaprozin as the next most effective followed by fenoprofen, naproxen, and ketoprofen as the least effective. Of these four drugs, naproxen has also been shown to suppress growth of prostate cancer cells; however, it was less effective when compared with ibuprofen (21).

The results showed that an induction of p75<sup>NTR</sup> was associated with significantly decreased survival of PC-3, DU-145, and LNCaP cells. Although these cell lines exhibited low levels of p75<sup>NTR</sup> expression at lower concentrations of *R*-flurbiprofen and ibuprofen, the levels of p75<sup>NTR</sup> were below the threshold for decreased survival. As the concentration of *R*-flurbiprofen or ibuprofen increased, some variability between levels of p75<sup>NTR</sup> and decreased




survival may have resulted from discordance around inflection points. At higher concentrations of *R*-flurbiprofen and ibuprofen, levels of p75<sup>NTR</sup> were concordant with decreased survival. This cause and effect relationship between higher levels of p75<sup>NTR</sup> and decreased survival was subsequently shown in the rescue experiments with dominant negatives and siRNA to p75<sup>NTR</sup>.

The exact mechanism by which *R*-flurbiprofen and ibuprofen induce p75<sup>NTR</sup> protein expression remains under investigation. However, an examination of p75<sup>NTR</sup> mRNA levels by RT-PCR at various time points showed that treatment of PC-3 and DU-145 cells with 2.0 mmol/L *R*-flurbiprofen or ibuprofen resulted in a relatively strong induction of p75<sup>NTR</sup> mRNA in all cases within 4 h and continuing until between 8 and 12 h, after which p75<sup>NTR</sup> mRNA remained constant. These results suggest that the observed induction of p75<sup>NTR</sup> protein expression is a result of an increase in p75<sup>NTR</sup> mRNA. This may occur by an increase in mRNA stability because previous work showed that p75<sup>NTR</sup> protein expression is lost in prostate cancer cell lines due to increased mRNA instability

that is mediated through the 3' untranslated region (31). Therefore, it is possible that treatment with *R*-flurbiprofen or ibuprofen alleviates this instability.

COX-2 catalyzes the conversion of arachidonic acid to prostaglandins, which are associated with increased survival, decreased apoptosis, and promotion of angiogenesis (13). Hence, inhibition of COX-2 has been suggested as a mechanism by which NSAIDs decrease cancer cell survival. However, an increasing body of literature suggests that many NSAIDs act through a COX-independent mechanism to achieve anticancer activity in the prostate (13). To elucidate whether COX plays a role in aryl propionic acid-dependent decreased prostate cancer cell survival, we examined the status of COX-1 and COX-2 expression. As expected, the housekeeping isoform COX-1 was expressed at low levels in PC-3 and DU-145 cells, and neither *R*-flurbiprofen nor ibuprofen induced its expression. Because COX-1 is not overexpressed in prostate cancer, is not generally associated with tumor development, and was minimally expressed in both cell lines at a



**Figure 6.** A to D, PC-3 and DU-145 cell survival analysis by MTT assay following 48-h treatment with 0, 0.5, or 1.0 mmol/L *R*-flurbiprofen or ibuprofen. Before treatment, cells were transfected with 100 nmol/L nontargeting siRNA or p75<sup>NTR</sup>-targeting siRNA for 24 h. Columns, mean relative to the control (0 mM); bars, SE. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ . p75<sup>NTR</sup> expression following 24-h transfection with 100 nmol/L nontargeting or p75<sup>NTR</sup>-targeting siRNA and subsequent 48-h treatment with 0, 0.5, or 1.0 mmol/L *R*-flurbiprofen or ibuprofen was determined by SDS-PAGE using 50  $\mu$ g of cell lysate and immunoblot analysis using a monoclonal antibody to human p75<sup>NTR</sup> and  $\beta$ -actin as the loading control.

constant level, it seems that *R*-flurbiprofen and ibuprofen inhibited survival through a mechanism other than COX-1 inhibition. DU-145 cells lacked any COX-2 expression, whereas PC-3 cells only exhibited expression upon *R*-flurbiprofen or ibuprofen treatment. An induction of COX-2 following NSAID treatment has been observed previously in PC-3 cells, as well as several other cell types (47–49). Although ibuprofen treatment led to an up-regulation of COX-2, the effect would be negated because ibuprofen would inhibit COX-2 activity. Because *R*-flurbiprofen lacks COX inhibitory activity, COX-2 could potentially remain active upon induction by *R*-flurbiprofen. However, 48-h *R*-flurbiprofen treatment of PC-3 cells was more potent for decreased survival than *R*-flurbiprofen in DU-145 cells, as well as more potent than ibuprofen treatment in both PC-3 and DU-145 cells. Therefore, there is no evidence that an induction of COX-2 in PC-3 cells promotes survival or reduces the effect of *R*-flurbiprofen treatment. Indeed, induction of COX-2 in PC-3 cells was associated with the greatest decrease in cell survival. This further supports the hypothesis that these drugs act as anticancer agents independently of COX-2.

p75<sup>NTR</sup> belongs to the TNFR superfamily, which includes Fas, p55<sup>TNFR</sup>, DR3, DR4, DR5, and DR6 (35). These receptors initiate instructive apoptosis through a homologous intracellular death domain (50). With the exception of p75<sup>NTR</sup>, they induce apoptosis upon ligand binding (51). In contrast, p75<sup>NTR</sup> initiates apoptosis and suppresses growth in a ligand-independent manner in prostate and bladder cancer cells (26–29). Previous studies show that some NSAIDs up-regulate expression of various death receptors (52, 53). To eliminate the possibility that the observed decrease in survival was due to apoptosis initiated by another member of the TNFR family, we examined expression of the remaining six family members in PC-3 cells following *R*-flurbiprofen or ibuprofen treatment. Neither p55<sup>TNFR</sup> nor DR6 were detected with or without treatment. Fas and DR5 were weakly expressed and remained constant upon treatment. DR3 expression was stronger than the other family members, but weak relative to p75<sup>NTR</sup>, and remained constant following treatment. DR4 seems slightly up-regulated by ibuprofen; however, its expression levels were minimal relative to the robust induction of p75<sup>NTR</sup>. Similar results were observed in the presence of the appropriate ligand for each receptor. Therefore, other than p75<sup>NTR</sup>, the death receptors capable of initiating apoptosis do not seem to play a significant role in the decreased survival caused by *R*-flurbiprofen and ibuprofen treatment of PC-3 and DU-145 prostate cancer cells.

It seems that *R*-flurbiprofen and ibuprofen decrease survival of PC-3 and DU-145 cells by a COX-independent mechanism. p75<sup>NTR</sup> is the only TNFR superfamily member significantly up-regulated by treatment with aryl propionic acids, and its up-regulation is associated with decreased survival in both cell lines. This suggests

p75<sup>NTR</sup> induction as a COX-independent mechanism initiating the observed decrease in survival. Hence, three different approaches including p75<sup>NTR</sup>-targeting siRNA and two different p75<sup>NTR</sup> dominant negatives were used to examine a causal relationship between p75<sup>NTR</sup> levels and decreased cell survival. Each method increased survival of PC-3 and DU-145 cells treated with *R*-flurbiprofen or ibuprofen relative to untransfected cells. Collectively, these results provide strong evidence that p75<sup>NTR</sup> is at least partially causal of the decreased survival due to treatment with *R*-flurbiprofen or ibuprofen.

Because chemopreventive and chemotherapeutic drugs are often administered chronically, we examined whether long-term treatment enabled *R*-flurbiprofen and ibuprofen to maintain their anticancer activity at lower concentrations than observed with acute treatments (48 h). Both *R*-flurbiprofen and ibuprofen were highly effective in inducing p75<sup>NTR</sup> and reducing growth at lower concentrations after chronic treatment. In particular, long-term *R*-flurbiprofen treatment at the lowest concentration of 0.1 mmol/L exhibited an 85% reduction of growth relative to the control and essentially total loss of growth at concentrations of 0.3 mmol/L or greater in PC-3 cells. Likewise, in DU-145 cells, 0.1 mmol/L long-term *R*-flurbiprofen treatment resulted in 66% reduction of growth and almost a total loss of growth at 0.5 mmol/L treatment. This dramatic loss of growth also correlates with the induction of p75<sup>NTR</sup>, which occurred in a dose-dependent manner starting at 0.1 mmol/L. Similarly, ibuprofen exhibited a substantial loss of growth due to chronic treatment at lower concentrations relative to acute 48-h treatment at higher concentrations in both PC-3 and DU-145 cells. The results of the chronic treatment experiments are especially significant because they indicate that these drugs, especially *R*-flurbiprofen, could be highly effective in inhibiting prostate cancer growth at clinically achievable concentrations with long-term use. In clinical trials, *R*-flurbiprofen and ibuprofen have been given at doses that result in plasma concentrations of 0.14 and 0.48 mmol/L, respectively (54). These concentrations are within the range of concentrations used in our chronic study. Hence, the data collectively show the activity of *R*-flurbiprofen and ibuprofen as anticancer agents in the prostate and convincingly implicates p75<sup>NTR</sup> induction as a COX-independent mechanism by which this anticancer activity is achieved.

## Acknowledgments

Received 10/3/2006; revised 12/29/2006; accepted 2/1/2007.

**Grant support:** NIH (R01DK52626), the Department of Defense Prostate Cancer Research Program (PC060409), and Myriad Pharmaceuticals Inc.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

## References

- Jemal A, Thomas A, Murray T, Thun M. Cancer statistics, 2002. *CA Cancer J Clin* 2002;52:23–47.
- Howe HL, Wingo PA, Thun MJ, et al. Annual report to the nation on the status of cancer (1973 through 1998), featuring cancers with recent increasing trends. *J Natl Cancer Inst* 2001;93:824–42.
- Garcia LA, Gonzalez-Perez A. Inverse association between nonsteroidal anti-inflammatory drugs and prostate cancer. *Cancer Epidemiol Biomarkers Prev* 2004;13:649–53.
- Roberts RO, Jacobson DJ, Girman CJ, Rhodes T, Lieber MM, Jacobsen SJ. A population-based study of daily

nonsteroidal anti-inflammatory drug use and prostate cancer. *Mayo Clin Proc* 2002;77:219–25.

- Nelson JE, Harris RE. Inverse association of prostate cancer and non-steroidal anti-inflammatory drugs (NSAIDs): results of a case-control study. *Oncol Rep* 2000;7:169–70.
- Habel LA, Zhao W, Stanford JL. Daily aspirin use and prostate cancer risk in a large, multiracial cohort in the US. *Cancer Causes Control* 2002;13:427–34.
- Liu XH, Yao S, Kirschenbaum A, Levine AC. NS398, a selective cyclooxygenase-2 inhibitor, induces apoptosis and down-regulates bcl-2 expression in LNCaP cells. *Cancer Res* 1998;58:4245–9.
- Gavrilov V, Steiner M, Shany S. The combined

treatment of 1,25-dihydroxyvitamin D3 and a non-steroid anti-inflammatory drug is highly effective in suppressing prostate cancer cell line (LNCaP) growth. *Anticancer Res* 2005;25:3425–9.

- Eberhart CE, Coffey RJ, Radhika A, Giardiello FM, Ferrenbach S, DuBois RN. Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology* 1994;107:1183–8.
- Gupta S, Srivastava M, Ahmad N, Bostwick DG, Mukhtar H. Over-expression of cyclooxygenase-2 in human prostate adenocarcinoma. *Prostate* 2000;42:73–8.
- Kirschenbaum A, Klausner AP, Lee R, et al. Expression of cyclooxygenase-1 and cyclooxygenase-2 in the human prostate. *Urology* 2000;56:671–6.

12. Madaan S, Abel PD, Chaudhary KS, et al. Cytoplasmic induction and over-expression of cyclooxygenase-2 in human prostate cancer: implications for prevention and treatment. *BJU Int* 2000;86:736-41.
13. Zha S, Gage WR, Sauvageot J, et al. Cyclooxygenase-2 is up-regulated in proliferative inflammatory atrophy of the prostate, but not in prostate carcinoma. *Cancer Res* 2001;61:8617-23.
14. Shappell SB, Manning S, Boeglin WE, et al. Alterations in lipoxygenase and cyclooxygenase-2 catalytic activity and mRNA expression in prostate carcinoma. *Neoplasia* 2001;3:287-303.
15. Subbarayan V, Sabichia AL, Llansa N, Lippman SM, Menter DG. Differential expression of cyclooxygenase-2 and its regulation by tumor necrosis factor- $\alpha$  in normal and malignant prostate cells. *Cancer Res* 2001;61:2720-6.
16. Lee JH, Cheong J, Park YM, Choi YH. Down-regulation of cyclooxygenase-2 and telomerase activity by  $\beta$ -lapachone in human prostate carcinoma cells. *Pharmacol Res* 2005;51:553-60.
17. Patel MI, Subbaramaiah K, Du B, et al. Celecoxib inhibits prostate cancer growth: evidence of a cyclooxygenase-2-independent mechanism. *Clin Cancer Res* 2005;11:1999-2007.
18. Song X, Lin HP, Johnson AJ, et al. Cyclooxygenase-2, player or spectator in cyclooxygenase-2 inhibitor-induced apoptosis in prostate cancer cells. *J Natl Cancer Inst* 2002;94:585-91.
19. Farivar-Mohseni H, Kandzari SJ, Zaslaw S, Riggs DR, Jackson BJ, McFadden DW. Synergistic effects of Cox-1 and -2 inhibition on bladder and prostate cancer *in vitro*. *Am J Surg* 2004;188:505-10.
20. Giovannucci E, Egan KM, Hunter DJ, et al. Aspirin and the risk of colorectal cancer in women. *N Engl J Med* 1995;333:609-14.
21. Andrews J, Djakiew D, Krygier S, Andrews P. Superior effectiveness of ibuprofen compared with other NSAIDs for reducing the survival of human prostate cancer cells. *Cancer Chemother Pharmacol* 2002;50:277-84.
22. Takada Y, Bhardwaj A, Potdar P, Aggarwal BB. Nonsteroidal anti-inflammatory agents differ in their ability to suppress NF- $\kappa$ B activation, inhibition of expression of cyclooxygenase-2 and cyclin D1, and abrogation of tumor cell proliferation. *Oncogene* 2004;23:9247-58.
23. Chao MV. The p75 neurotrophin receptor. *J Neurobiol* 1994;25:1373-85.
24. Chapman BS. A region of the 75 kDa neurotrophin receptor homologous to the death domains of TNFR-I and Fas. *FEBS Lett* 1995;374:216-20.
25. Friedman WJ, Greene LA. Neurotrophin signaling via Trks and p75. *Exp Cell Res* 1999;253:131-42.
26. Pflug B, Djakiew D. Expression of p75<sup>NTR</sup> in a human prostate epithelial tumor cell line reduces nerve growth factor-induced cell growth by activation of programmed cell death. *Mol Carcinog* 1998;23:106-14.
27. Krygier S, Djakiew D. The neurotrophin receptor p75<sup>NTR</sup> is a tumor suppressor in human prostate cancer. *Anticancer Res* 2001;21:3749-55.
28. Krygier S, Djakiew D. Neurotrophin receptor p75<sup>NTR</sup> suppresses growth and nerve growth factor-mediated metastasis of human prostate cancer cells. *Int J Cancer* 2002;98:1-7.
29. Khwaja F, Tabassum A, Allen J, Djakiew D. The p75<sup>NTR</sup> tumor suppressor induces cell cycle arrest facilitating caspase mediated apoptosis in prostate tumor cells. *Biochem Biophys Res Commun* 2006;341:1184-92.
30. Perez M, Regan T, Pflug B, Lynch J, Djakiew D. Loss of low-affinity nerve growth factor receptor during malignant transformation of the human prostate. *Prostate* 1997;30:274-9.
31. Krygier S, Djakiew D. Molecular characterization of the loss of p75<sup>NTR</sup> expression in human prostate tumor cells. *Mol Carcinog* 2001;31:46-55.
32. Allen J, Khwaja F, Byers S, Djakiew D. The p75<sup>NTR</sup> mediates a bifurcated signal transduction cascade through the NF- $\kappa$ B and JNK pathways to inhibit cell survival. *Exp Cell Res* 2005;304:69-80.
33. Khwaja F, Allen J, Lynch J, Andrews P, Djakiew D. Ibuprofen inhibits survival of bladder cancer cells by induced expression of the p75<sup>NTR</sup> tumor suppressor protein. *Cancer Res* 2004;64:6207-13.
34. Wechter WJ, Leipold DD, Murray ED, Jr., et al. E-7869 (R-flurbiprofen) inhibits progression of prostate cancer in the TRAMP mouse. *Cancer Res* 2000;60:2203-8.
35. Wang X, Bauer JH, Li Y, et al. Characterization of a p75<sup>NTR</sup> apoptotic signaling pathway using a novel cellular model. *J Biol Chem* 2001;276:33812-20.
36. Evans AM. Pharmacodynamics and pharmacokinetics of the profens: enantioselectivity, clinical implications, and special reference to S(+)-ibuprofen. *J Clin Pharmacol* 1996;36:7-15S.
37. Landoni MF, Soraci A. Pharmacology of chiral compounds: 2-arylpropionic acid derivatives. *Curr Drug Metab* 2001;2:37-51.
38. Rovinsky J, Micekova D. Six-month prospective study to monitor the treatment of rheumatic diseases with sustained-release flurbiprofen. *Drugs Exp Clin Res* 2000;26:19-24.
39. Kean WF. Oxaprozin: kinetic and dynamic profile in the treatment of pain. *Curr Med Res Opin* 2004;20:1275-7.
40. Grahame R. Ketoprofen-clinical efficacy. *Rheumatol Rehabil* 1976;Suppl:22-6.
41. Ward JR. Update on ibuprofen for rheumatoid arthritis. *Am J Med* 1984;77:3-9.
42. Sigler JW, Ridolfo AS, Bluhm GB. Comparison of benefit-to-risk ratios of aspirin and fenoprofen: controlled multicentre study in rheumatoid arthritis. *J Rheumatol* 1976;2:49-60.
43. Lee P, Rose BS, Anderson JA, Caughey DE. Naproxen in the treatment of rheumatoid arthritis. *N Z Med J* 1978;87:425-7.
44. Wargovich MJ, Jimenez A, McKee K, et al. Efficacy of potential chemopreventive agents on rat colon aberrant crypt formation and progression. *Carcinogenesis* 2000;21:1149-55.
45. Harris RE, Chlebowski RT, Jackson RD, et al. Breast cancer and nonsteroidal anti-inflammatory drugs: prospective results from the Women's Health Initiative. *Cancer Res* 2003;63:6096-101.
46. Castonguay A, Rioux N, Duperron C, Jalbert G. Inhibition of lung tumorigenesis by NSAIDs: a working hypothesis. *Exp Lung Res* 1998;24:605-15.
47. Meade EA, McIntyre TM, Zimmerman GA, Prescott SM. Peroxisome proliferators enhance cyclooxygenase-2 expression in epithelial cells. *J Biol Chem* 1999;274:8328-34.
48. Yuan CJ, Mandal AK, Zhang Z, Mukherjee AB. Transcriptional regulation of cyclooxygenase-2 gene expression: novel effects of nonsteroidal anti-inflammatory drugs. *Cancer Res* 2000;60:1084-91.
49. Palayoor ST, Tofilon PJ, Coleman CN. Ibuprofen-mediated reduction of hypoxia-inducible factors HIF-1 $\alpha$  and HIF-2 $\alpha$  in prostate cancer cells. *Clin Cancer Res* 2003;9:3150-7.
50. Ashkenazi A, Dixit VM. Apoptosis control by death and decoy receptors. *Curr Opin Cell Biol* 1999;11:255-60.
51. Ashkenazi A, Dixit VM. Death receptors: signaling and modulation. *Science* 1998;281:1305-8.
52. Zhang M, Abe Y, Matsushima T, Nishimura J, Nawata H, Muta K. Selective cyclooxygenase 2 inhibitor NS-398 induces apoptosis in myeloma cells via a Bcl-2 independent pathway. *Leuk Lymphoma* 2005;46:425-33.
53. Huang Y, He Q, Hillman MJ, Rong R, Sheikh MS. Sulindac sulfide-induced apoptosis involves death receptor 5 and the caspase 8-dependent pathway in human colon and prostate cancer cells. *Cancer Res* 2001;61:6918-24.
54. Konstan MW, Byard PJ, Hoppel CL, Davis PB. Effect of high-dose ibuprofen in patients with cystic fibrosis. *N Engl J Med* 1995;332:848-54.